



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 15, 2020 – 10:29 pm BST

PDB ID : 1B6G
Title : HALOALKANE DEHALOGENASE AT PH 5.0 CONTAINING CHLORIDE
Authors : Ridder, I.S.; Rozeboom, H.J.; Dijkstra, B.W.
Deposited on : 1999-01-14
Resolution : 1.15 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	1537 (1.18-1.10)
Ramachandran outliers	138981	1483 (1.18-1.10)
Sidechain outliers	138945	1480 (1.18-1.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	310	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	GOL	A	1202	-	X	-	-
4	GOL	A	1204	-	-	X	-

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 3289 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called HALOALKANE DEHALOGENASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	N	O	Pb				S
1	A	310	2639	1694	429	497	2	17	0	34	0

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	2	VAL	ILE	CLONING ARTIFACT	UNP P22643
A	150	CSB	CYS	MODIFIED RESIDUE	UNP P22643
A	233	CSB	CYS	MODIFIED RESIDUE	UNP P22643

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O S		
2	A	1	5	4 1	0	0

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	2	Total Cl 2 2	0	0

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 6 3 3	0	0
4	A	1	Total C O 6 3 3	0	0
4	A	1	Total C O 6 3 3	0	0
4	A	1	Total C O 6 3 3	0	0
4	A	1	Total C O 6 3 3	0	0
4	A	1	Total C O 6 3 3	0	0
4	A	1	Total C O 6 3 3	0	0

- Molecule 5 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	601	Total O 601 601	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: HALOALKANE DEHALOGENASE



4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	92.18Å 72.03Å 40.91Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	15.00 – 1.15	Depositor
% Data completeness (in resolution range)	97.6 (15.00-1.15)	Depositor
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	SHELXL-97	Depositor
R, R_{free}	0.105 , 0.145	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	3289	wwPDB-VP
Average B, all atoms (Å ²)	18.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, CL, SO4, CSB

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.65	1/2816 (0.0%)	1.14	19/3820 (0.5%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	56	GLU	CD-OE2	5.30	1.31	1.25

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	16	ASP	CB-CG-OD1	7.64	125.18	118.30
1	A	9[A]	ASP	CB-CG-OD2	-7.22	111.80	118.30
1	A	9[B]	ASP	CB-CG-OD2	-7.22	111.80	118.30
1	A	228[A]	GLN	CB-CA-C	6.36	123.11	110.40
1	A	228[B]	GLN	CB-CA-C	6.36	123.11	110.40
1	A	225[A]	MET	CA-CB-CG	-6.00	103.11	113.30
1	A	225[B]	MET	CA-CB-CG	-6.00	103.11	113.30
1	A	31	TYR	CG-CD2-CE2	5.52	125.72	121.30
1	A	186[A]	ARG	CD-NE-CZ	5.49	131.29	123.60
1	A	186[B]	ARG	CD-NE-CZ	5.49	131.29	123.60
1	A	63	TYR	CB-CG-CD2	5.41	124.25	121.00
1	A	11	ARG	NE-CZ-NH1	5.40	123.00	120.30
1	A	63	TYR	CB-CG-CD1	-5.36	117.78	121.00
1	A	27	ASP	CB-CG-OD1	5.36	123.12	118.30
1	A	31	TYR	CA-CB-CG	-5.29	103.35	113.40
1	A	18	TYR	CB-CG-CD2	5.27	124.16	121.00
1	A	306	PHE	CB-CG-CD1	5.10	124.37	120.80
1	A	143[A]	ARG	CD-NE-CZ	5.01	130.62	123.60
1	A	143[B]	ARG	CD-NE-CZ	5.01	130.62	123.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2639	0	2515	33	0
2	A	5	0	0	1	0
3	A	2	0	0	0	0
4	A	42	0	53	8	0
5	A	601	0	0	31	0
All	All	3289	0	2568	40	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (40) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:181[B]:THR:HG22	5:A:2514:HOH:O	1.47	1.15
1:A:213[C]:THR:OG1	5:A:2578:HOH:O	1.66	1.12
1:A:192[B]:LYS:HD3	5:A:2551:HOH:O	0.91	1.08
1:A:9[B]:ASP:OD1	5:A:2598:HOH:O	1.70	1.07
2:A:1101:SO4:O4	5:A:2581:HOH:O	1.74	1.05
4:A:1204:GOL:H2	5:A:2609:HOH:O	0.89	1.05
1:A:181[B]:THR:CG2	5:A:2514:HOH:O	2.04	1.04
1:A:213[A]:THR:HG22	5:A:2578:HOH:O	0.82	0.98
1:A:137[A]:ASP:HB2	5:A:2585:HOH:O	1.81	0.81
1:A:213[B]:THR:HG22	5:A:2455:HOH:O	1.83	0.78
4:A:1205:GOL:H12	5:A:2339:HOH:O	1.86	0.75
1:A:310:GLU:HG2	5:A:2217:HOH:O	1.89	0.72
1:A:192[B]:LYS:NZ	5:A:2551:HOH:O	2.18	0.72
1:A:213[A]:THR:CG2	5:A:2578:HOH:O	1.64	0.71
1:A:192[B]:LYS:CD	5:A:2551:HOH:O	1.71	0.70
1:A:95[A]:GLU:OE2	4:A:1204:GOL:C1	2.40	0.69
1:A:306:PHE:O	1:A:310:GLU:HG3	1.94	0.68
1:A:176[B]:LYS:NZ	5:A:2509:HOH:O	1.83	0.67

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:9[A]:ASP:OD2	5:A:2597:HOH:O	2.13	0.66
1:A:1:MET:SD	1:A:33:GLY:HA3	2.45	0.57
1:A:9[A]:ASP:OD1	5:A:2590:HOH:O	2.17	0.57
4:A:1204:GOL:H32	5:A:2103:HOH:O	2.03	0.56
1:A:47[B]:GLU:O	1:A:48[B]:ASP:OD1	2.25	0.54
1:A:111[B]:GLU:HG2	5:A:2250:HOH:O	2.08	0.54
1:A:11:ARG:NH2	5:A:2577:HOH:O	2.41	0.54
1:A:213[A]:THR:HG23	5:A:2205:HOH:O	2.09	0.51
1:A:183:SER:O	1:A:213[A]:THR:HG21	2.11	0.50
1:A:111[A]:GLU:HG3	5:A:2250:HOH:O	2.12	0.49
1:A:239[A]:GLU:HG3	5:A:2416:HOH:O	2.11	0.49
1:A:181[B]:THR:HG21	5:A:2514:HOH:O	1.89	0.48
1:A:192[B]:LYS:NZ	5:A:2369:HOH:O	2.46	0.47
4:A:1204:GOL:C3	5:A:2609:HOH:O	2.29	0.46
1:A:187:LEU:HD12	1:A:210:PHE:CD2	2.49	0.46
4:A:1207:GOL:H2	5:A:2069:HOH:O	2.16	0.46
1:A:228[A]:GLN:NE2	5:A:2267:HOH:O	2.49	0.44
1:A:152[B]:MET:HG3	5:A:2064:HOH:O	2.18	0.44
1:A:1:MET:HB2	1:A:32:PRO:O	2.20	0.42
4:A:1204:GOL:C2	5:A:2609:HOH:O	1.79	0.42
1:A:125:TRP:CH2	1:A:152[B]:MET:SD	3.14	0.41
1:A:294:PHE:O	4:A:1206:GOL:O2	2.39	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	339/310 (109%)	330 (97%)	8 (2%)	1 (0%)	41 16

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	148	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	293/260 (113%)	289 (99%)	4 (1%)	67 29

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	3	ASN
1	A	72[A]	GLU
1	A	72[B]	GLU
1	A	222	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	3	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 2 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	GOL	A	1203	-	5,5,5	0.66	0	5,5,5	1.66	2 (40%)
4	GOL	A	1207	-	5,5,5	0.47	0	5,5,5	1.35	1 (20%)
4	GOL	A	1204	1	5,5,5	0.72	0	5,5,5	1.72	1 (20%)
4	GOL	A	1201	-	5,5,5	0.51	0	5,5,5	0.72	0
2	SO4	A	1101	1	4,4,4	0.51	0	6,6,6	0.37	0
4	GOL	A	1205	-	5,5,5	0.43	0	5,5,5	0.84	0
4	GOL	A	1202	-	5,5,5	0.75	0	5,5,5	2.20	3 (60%)
4	GOL	A	1206	-	5,5,5	0.84	0	5,5,5	1.42	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	A	1203	-	-	2/4/4/4	-
4	GOL	A	1207	-	-	1/4/4/4	-
4	GOL	A	1204	1	-	3/4/4/4	-
4	GOL	A	1201	-	-	0/4/4/4	-
4	GOL	A	1205	-	-	3/4/4/4	-
4	GOL	A	1202	-	-	4/4/4/4	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	A	1206	-	-	3/4/4/4	-

There are no bond length outliers.

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	1204	GOL	O2-C2-C3	3.66	125.25	109.12
4	A	1202	GOL	O2-C2-C1	2.92	121.99	109.12
4	A	1202	GOL	O2-C2-C3	-2.82	96.71	109.12
4	A	1207	GOL	O2-C2-C3	2.77	121.32	109.12
4	A	1203	GOL	O2-C2-C1	2.57	120.44	109.12
4	A	1202	GOL	O3-C3-C2	2.56	122.47	110.20
4	A	1203	GOL	C3-C2-C1	2.33	120.76	111.70

There are no chirality outliers.

All (16) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	1202	GOL	O1-C1-C2-C3
4	A	1206	GOL	O1-C1-C2-C3
4	A	1204	GOL	O2-C2-C3-O3
4	A	1203	GOL	C1-C2-C3-O3
4	A	1202	GOL	C1-C2-C3-O3
4	A	1204	GOL	O1-C1-C2-C3
4	A	1207	GOL	O1-C1-C2-C3
4	A	1205	GOL	C1-C2-C3-O3
4	A	1202	GOL	O1-C1-C2-O2
4	A	1202	GOL	O2-C2-C3-O3
4	A	1205	GOL	O1-C1-C2-O2
4	A	1206	GOL	O1-C1-C2-O2
4	A	1203	GOL	O1-C1-C2-O2
4	A	1206	GOL	O2-C2-C3-O3
4	A	1204	GOL	C1-C2-C3-O3
4	A	1205	GOL	O1-C1-C2-C3

There are no ring outliers.

5 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	1207	GOL	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	1204	GOL	5	0
2	A	1101	SO4	1	0
4	A	1205	GOL	1	0
4	A	1206	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.