



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 2, 2021 – 04:16 AM EDT

PDB ID : 2FXO
Title : Structure of the human beta-myosin S2 fragment
Authors : Blankenfeldt, W.; Thoma, N.H.; Wray, J.S.; Gautel, M.; Schlichting, I.
Deposited on : 2006-02-06
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.23.2
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.23.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4661 (2.50-2.50)
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	129	 3% (poor fit), 71% (0-1 outliers), 26% (2-3 outliers), . (not modelled)
1	B	129	 75% (0-1 outliers), 21% (2-3 outliers), .. (not modelled)
1	C	129	 2% (poor fit), 74% (0-1 outliers), 24% (2-3 outliers), . (not modelled)
1	D	129	 72% (0-1 outliers), 24% (2-3 outliers), .. (not modelled)

2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 4145 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Myosin heavy chain, cardiac muscle beta isoform.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	129	1048	637	183	221	7	17	0	0
1	B	125	1024	623	178	216	7	16	0	0
1	C	127	1035	629	180	219	7	4	0	0
1	D	127	1038	632	181	218	7	35	0	0


There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-3	GLY	-	cloning artifact	UNP P12883
A	-2	SER	-	cloning artifact	UNP P12883
A	-1	SER	-	cloning artifact	UNP P12883
A	924	LYS	GLU	engineered mutation	UNP P12883
B	-3	GLY	-	cloning artifact	UNP P12883
B	-2	SER	-	cloning artifact	UNP P12883
B	-1	SER	-	cloning artifact	UNP P12883
B	924	LYS	GLU	engineered mutation	UNP P12883
C	-3	GLY	-	cloning artifact	UNP P12883
C	-2	SER	-	cloning artifact	UNP P12883
C	-1	SER	-	cloning artifact	UNP P12883
C	924	LYS	GLU	engineered mutation	UNP P12883
D	-3	GLY	-	cloning artifact	UNP P12883
D	-2	SER	-	cloning artifact	UNP P12883
D	-1	SER	-	cloning artifact	UNP P12883
D	924	LYS	GLU	engineered mutation	UNP P12883


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Myosin heavy chain, cardiac muscle beta isoform




- Molecule 1: Myosin heavy chain, cardiac muscle beta isoform



- Molecule 1: Myosin heavy chain, cardiac muscle beta isoform



- Molecule 1: Myosin heavy chain, cardiac muscle beta isoform



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	40.16Å 41.87Å 97.80Å 91.11° 92.73° 107.18°	Depositor
Resolution (Å)	20.00 – 2.50 97.62 – 2.40	Depositor EDS
% Data completeness (in resolution range)	96.2 (20.00-2.50) 96.2 (97.62-2.40)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.31 (at 2.40Å)	Xtrriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.273 , 0.349 0.269 , 0.346	Depositor DCC
R_{free} test set	1040 reflections (4.38%)	wwPDB-VP
Wilson B-factor (Å ²)	56.1	Xtrriage
Anisotropy	0.223	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 50.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.52$, $\langle L^2 \rangle = 0.35$	Xtrriage
Estimated twinning fraction	0.000 for -k,-h,-l	Xtrriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	4145	wwPDB-VP
Average B, all atoms (Å ²)	71.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 13.72% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.67	0/1049	0.90	5/1391 (0.4%)
1	B	0.71	0/1025	0.91	5/1360 (0.4%)
1	C	0.69	1/1036 (0.1%)	0.88	4/1375 (0.3%)
1	D	0.67	0/1039	0.90	6/1378 (0.4%)
All	All	0.69	1/4149 (0.0%)	0.90	20/5504 (0.4%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	947	CYS	CB-SG	5.87	1.92	1.82

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	906	ASP	CB-CG-OD2	7.52	125.06	118.30
1	D	955	ASP	CB-CG-OD2	6.63	124.27	118.30
1	D	953	ASP	CB-CG-OD2	6.11	123.80	118.30
1	D	906	ASP	CB-CG-OD2	5.98	123.69	118.30
1	A	945	ASP	CB-CG-OD2	5.87	123.58	118.30
1	D	945	ASP	CB-CG-OD2	5.87	123.58	118.30
1	A	906	ASP	CB-CG-OD2	5.86	123.57	118.30
1	B	900	ASP	CB-CG-OD2	5.77	123.50	118.30
1	A	896	ASP	CB-CG-OD2	5.67	123.40	118.30
1	C	896	ASP	CB-CG-OD2	5.67	123.40	118.30
1	D	900	ASP	CB-CG-OD2	5.51	123.26	118.30
1	A	955	ASP	CB-CG-OD2	5.49	123.24	118.30
1	B	845	ARG	NE-CZ-NH1	5.41	123.00	120.30
1	C	956	ASP	CB-CG-OD2	5.33	123.09	118.30
1	B	906	ASP	CB-CG-OD2	5.32	123.08	118.30
1	D	928	ASP	CB-CG-OD2	5.30	123.07	118.30
1	B	955	ASP	CB-CG-OD2	5.25	123.02	118.30
1	B	896	ASP	CB-CG-OD2	5.12	122.91	118.30
1	A	928	ASP	CB-CG-OD2	5.10	122.89	118.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	945	ASP	CB-CG-OD2	5.09	122.88	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1048	0	1071	12	0
1	B	1024	0	1045	8	0
1	C	1035	0	1055	8	0
1	D	1038	0	1063	12	0
All	All	4145	0	4234	29	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (29) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:907:GLN:OE1	1:D:865:LYS:NZ	2.13	0.82
1:A:933:ASN:ND2	1:B:933:ASN:OD1	2.13	0.81
1:D:918:LYS:O	1:D:922:MET:HG2	1.82	0.79
1:A:874:GLU:OE2	1:B:869:ARG:NE	2.21	0.71
1:B:884:LYS:HE2	1:B:888:GLN:HE22	1.57	0.69
1:B:884:LYS:CE	1:B:888:GLN:HE22	2.10	0.64
1:D:867:GLU:OE1	1:D:870:ARG:NH1	2.32	0.63
1:A:891:VAL:O	1:A:895:GLN:HG3	2.02	0.59
1:D:-1:SER:N	1:D:838:PRO:CD	2.68	0.57
1:A:891:VAL:HG23	1:B:891:VAL:HG23	1.86	0.56
1:D:-1:SER:H3	1:D:838:PRO:HD3	1.77	0.50
1:C:880:LEU:HD23	1:D:880:LEU:HD23	1.94	0.48
1:A:886:ASP:OD1	1:A:886:ASP:C	2.53	0.47
1:A:887:LEU:HA	1:A:890:GLN:HB2	1.97	0.47
1:C:-2:SER:HB2	1:C:839:LEU:HD12	1.97	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:898:LEU:HD13	1:D:897:ASN:HB3	1.96	0.46
1:C:-2:SER:CB	1:C:839:LEU:HD12	2.46	0.46
1:A:922:MET:HE1	1:B:919:VAL:HG13	2.00	0.43
1:C:891:VAL:HG23	1:D:891:VAL:HG23	2.00	0.43
1:C:880:LEU:HD21	1:D:881:LEU:HG	2.00	0.43
1:A:-1:SER:HB3	1:A:838:PRO:HD3	2.00	0.42
1:D:-1:SER:H3	1:D:838:PRO:CD	2.33	0.41
1:A:911:ASN:HD21	1:B:912:LYS:HE3	1.86	0.41
1:C:926:LEU:HG	1:D:926:LEU:HD12	2.01	0.41
1:A:924:LYS:HG2	1:A:925:ARG:N	2.36	0.41
1:A:869:ARG:O	1:A:873:LEU:HB2	2.21	0.40
1:D:887:LEU:O	1:D:888:GLN:C	2.60	0.40
1:A:849:MET:O	1:A:853:LYS:HG2	2.21	0.40
1:C:937:THR:O	1:C:938:ALA:C	2.59	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	127/129 (98%)	124 (98%)	3 (2%)	0	100	100
1	B	123/129 (95%)	117 (95%)	6 (5%)	0	100	100
1	C	125/129 (97%)	120 (96%)	5 (4%)	0	100	100
1	D	125/129 (97%)	116 (93%)	8 (6%)	1 (1%)	19	35
All	All	500/516 (97%)	477 (95%)	22 (4%)	1 (0%)	47	68

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	962	ALA

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	117/117 (100%)	99 (85%)	18 (15%)	2	5
1	B	115/117 (98%)	99 (86%)	16 (14%)	3	6
1	C	116/117 (99%)	98 (84%)	18 (16%)	2	4
1	D	116/117 (99%)	100 (86%)	16 (14%)	3	6
All	All	464/468 (99%)	396 (85%)	68 (15%)	3	5

All (68) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	-1	SER
1	A	840	LEU
1	A	841	LYS
1	A	855	GLU
1	A	856	PHE
1	A	857	THR
1	A	858	ARG
1	A	865	LYS
1	A	866	SER
1	A	867	GLU
1	A	874	GLU
1	A	875	GLU
1	A	877	MET
1	A	879	SER
1	A	880	LEU
1	A	881	LEU
1	A	887	LEU
1	A	935	GLU
1	B	839	LEU
1	B	845	ARG
1	B	847	LYS
1	B	848	GLU
1	B	851	SER
1	B	853	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	858	ARG
1	B	863	LEU
1	B	864	GLU
1	B	872	GLU
1	B	879	SER
1	B	881	LEU
1	B	921	GLU
1	B	926	LEU
1	B	927	GLU
1	B	928	ASP
1	C	-1	SER
1	C	844	GLU
1	C	849	MET
1	C	851	SER
1	C	858	ARG
1	C	866	SER
1	C	875	GLU
1	C	884	LYS
1	C	903	GLU
1	C	910	LYS
1	C	912	LYS
1	C	913	ILE
1	C	918	LYS
1	C	922	MET
1	C	927	GLU
1	C	933	ASN
1	C	957	LEU
1	C	959	LEU
1	D	847	LYS
1	D	849	MET
1	D	853	LYS
1	D	861	GLU
1	D	864	GLU
1	D	869	ARG
1	D	874	GLU
1	D	883	GLU
1	D	897	ASN
1	D	900	ASP
1	D	906	ASP
1	D	907	GLN
1	D	911	ASN
1	D	920	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	D	924	LYS
1	D	932	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	892	GLN
1	A	895	GLN
1	A	933	ASN
1	B	888	GLN
1	C	933	ASN
1	D	885	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	129/129 (100%)	-0.18	4 (3%) 49 52	35, 65, 92, 107	5 (3%)
1	B	125/129 (96%)	-0.24	0 100 100	31, 58, 90, 96	5 (4%)
1	C	127/129 (98%)	-0.10	2 (1%) 72 74	40, 63, 80, 90	1 (0%)
1	D	127/129 (98%)	-0.15	0 100 100	39, 68, 85, 91	11 (8%)
All	All	508/516 (98%)	-0.17	6 (1%) 79 80	31, 65, 87, 107	22 (4%)

All (6) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	-3	GLY	6.0
1	C	962	ALA	4.4
1	A	962	ALA	2.6
1	A	-2	SER	2.5
1	C	-2	SER	2.1
1	A	864	GLU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.