



## wwPDB/EMDatabank EM Map/Model Validation Summary Report ⓘ

Sep 19, 2019 – 10:22 AM EDT

PDB ID : 6QX3  
EMDB ID: : EMD-4661  
Title : Influenza A virus (A/NT/60/1968) polymerase Heterotrimer in complex with  
3'5' cRNA promoter and Nb8205  
Authors : Carrique, L.; Keown, J.R.; Fan, H.; Fodor, E.; Grimes, J.M.  
Deposited on : 2019-03-07  
Resolution : 3.79 Å(reported)  
Based on PDB ID : 6QPG

This is a wwPDB/EMDatabank EM Map/Model Validation Summary Report  
for a publicly released PDB/EMDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---


MolProbity : 4.02b-467  
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.4

# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:  
*ELECTRON MICROSCOPY*

The reported resolution of this entry is 3.79 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	136327	1886
Ramachandran outliers	132723	1663
Sidechain outliers	132532	1531
RNA backbone	3747	458

The table below summarises the geometric issues observed across the polymeric chains. The red, orange, yellow and green segments on the bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ .

Mol	Chain	Length	Quality of chain
1	D	15	33% (green), 27% (yellow), 7% (orange), 33% (grey)
2	G	15	7% (yellow), 13% (orange), 7% (red), 73% (grey)
3	O	134	83% (green), 7% (yellow), 10% (grey)
4	B	757	73% (green), 9% (yellow), 19% (grey)
5	A	716	64% (green), 8% (yellow), 28% (grey)
6	C	762	10% (green), 89% (grey)

## 2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 21587 atoms, of which 10646 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a RNA chain called RNA (5'-R(P\*AP\*GP\*CP\*AP\*AP\*AP\*AP\*GP\*CP\*A)-3').

Mol	Chain	Residues	Atoms						AltConf	Trace
			Total	C	H	N	O	P		
1	D	10	329	98	111	46	64	10	0	0

- Molecule 2 is a RNA chain called RNA (5'-R(P\*UP\*UP\*CP\*U)-3').

Mol	Chain	Residues	Atoms						AltConf	Trace
			Total	C	H	N	O	P		
2	G	4	122	36	42	9	31	4	0	0

- Molecule 3 is a protein called Nb8205.

Mol	Chain	Residues	Atoms						AltConf	Trace
			Total	C	H	N	O	S		
3	O	121	1799	572	877	163	181	6	0	0

- Molecule 4 is a protein called RNA-directed RNA polymerase catalytic subunit.

Mol	Chain	Residues	Atoms						AltConf	Trace
			Total	C	H	N	O	S		
4	B	615	9750	3084	4858	850	923	35	0	0

- Molecule 5 is a protein called Polymerase acidic protein.

Mol	Chain	Residues	Atoms						AltConf	Trace
			Total	C	H	N	O	S		
5	A	515	8157	2614	4039	691	783	30	0	0

- Molecule 6 is a protein called Polymerase basic protein 2.

Mol	Chain	Residues	Atoms						AltConf	Trace
			Total	C	H	N	O	S		
6	C	86	1430	453	719	125	125	8	0	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	757	GLU	-	expression tag	UNP P03429
C	758	ASN	-	expression tag	UNP P03429
C	759	LEU	-	expression tag	UNP P03429
C	760	TYR	-	expression tag	UNP P03429
C	761	PHE	-	expression tag	UNP P03429
C	762	GLN	-	expression tag	UNP P03429





ARG  
LEU  
THR  
ILE  
LEU  
GLY  
GLY  
LYS  
ASP  
ALA  
GLY  
THR  
LEU  
ILE  
GLU  
ASP  
PRO  
ASP  
GLY  
THR  
SER  
GLY  
VAL  
GLU  
SER  
ALA  
VAL  
LEU  
ARG  
GLY  
PHE  
LEU  
ILE  
LEU  
GLY  
LYS  
GLU  
ASP  
ARG  
TYR  
GLY  
PRO  
ALA  
LEU  
SER  
ILE  
ASN  
GLU  
LEU  
SER  
ASN  
ALA  
LYS  
GLY  
GLU  
LYS  
ALA  
ASN

VAL  
LEU  
ILE  
GLY  
GLN  
GLY  
ASP  
VAL  
VAL  
LEU  
VAL  
MET  
LYS  
ARG  
LYS  
ARG  
ASP  
SER  
SER  
ILE  
LEU  
THR  
ASP  
SER  
GLN  
THR  
ALA  
THR  
LYS  
ARG  
ILE  
ARG  
MET  
GLU  
ASN  
LEU  
TYR  
PHE  
GLN

## 4 Experimental information

Property	Value	Source
Reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C1	Depositor
Number of particles used	52932	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ( $e^-/\text{\AA}^2$ )	1.25	Depositor
Minimum defocus (nm)	1300	Depositor
Maximum defocus (nm)	2500	Depositor
Magnification	130000	Depositor
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >2	RMSZ	# Z  >2
1	D	0.29	0/245	0.72	0/380
2	G	0.36	0/87	1.68	5/132 (3.8%)
3	O	0.35	0/942	0.56	0/1278
4	B	0.33	0/4991	0.54	0/6744
5	A	0.37	0/4209	0.54	0/5688
6	C	0.27	0/730	0.51	0/985
All	All	0.34	0/11204	0.56	5/15207 (0.0%)

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	G	11	C	C2-N1-C1'	9.14	128.85	118.80
2	G	11	C	N1-C2-O2	7.03	123.12	118.90
2	G	11	C	C6-N1-C1'	-6.33	113.20	120.80
2	G	11	C	N3-C2-O2	-6.18	117.57	121.90
2	G	11	C	C6-N1-C2	-5.93	117.93	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	D	218	111	111	3	0
2	G	80	42	42	6	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	O	922	877	877	7	0
4	B	4892	4858	4856	42	0
5	A	4118	4039	4037	39	0
6	C	711	719	719	7	0
All	All	10941	10646	10642	86	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 86 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:B:563:ARG:O	4:B:566:THR:OG1	2.05	0.74
5:A:566:ARG:NE	5:A:567:THR:O	2.22	0.72
3:O:90:ASP:OD1	3:O:94:TYR:OH	2.08	0.71
4:B:619:ASP:OD1	4:B:623:ARG:NH1	2.25	0.70
2:G:11:C:OP2	5:A:488:LYS:NZ	2.24	0.69

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	O	119/134 (89%)	108 (91%)	11 (9%)	0	100	100
4	B	609/757 (80%)	549 (90%)	60 (10%)	0	100	100
5	A	513/716 (72%)	463 (90%)	50 (10%)	0	100	100
6	C	84/762 (11%)	77 (92%)	7 (8%)	0	100	100
All	All	1325/2369 (56%)	1197 (90%)	128 (10%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	O	97/111 (87%)	97 (100%)	0	100	100
4	B	538/669 (80%)	538 (100%)	0	100	100
5	A	453/644 (70%)	450 (99%)	3 (1%)	85	93
6	C	79/674 (12%)	79 (100%)	0	100	100
All	All	1167/2098 (56%)	1164 (100%)	3 (0%)	93	97

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
5	A	496	ARG
5	A	506	LYS
5	A	551	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
4	B	629	ASN
5	A	587	GLN

### 5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
1	D	9/15 (60%)	3 (33%)	0
2	G	3/15 (20%)	3 (100%)	0
All	All	12/30 (40%)	6 (50%)	0

5 of 6 RNA backbone outliers are listed below:

Mol	Chain	Res	Type
1	D	5	A
1	D	6	A

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	D	10	A
2	G	10	U
2	G	11	C

There are no RNA pucker outliers to report.

#### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

#### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

#### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

#### 5.7 Other polymers [i](#)

There are no such residues in this entry.

#### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.