



wwPDB EM Validation Summary Report ⓘ

Dec 6, 2020 – 05:28 AM GMT

PDB ID : 6R93
EMDB ID : EMD-4767
Title : Cryo-EM structure of NCP-6-4PP
Authors : Matsumoto, S.; Cavadini, S.; Bunker, R.D.; Thoma, N.H.
Deposited on : 2019-04-02
Resolution : 4.00 Å(reported)
Based on initial models : 4ZUX, 4E54, 3EI4, 5Y0C

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


EMDB validation analysis : 0.0.0.dev61
Mogul : 1.8.5 (274361), CSD as541be (2020)
MolProbity : 4.02b-467
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.15.1

1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 4.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	I	147	15% (red), 89% (green), 9% (yellow), 0% (orange), 0% (grey)
2	J	147	12% (red), 90% (green), 10% (yellow), 0% (orange), 0% (grey)
3	A	139	19% (red), 63% (green), 8% (yellow), 0% (orange), 29% (grey)
3	E	139	22% (red), 64% (green), 7% (yellow), 0% (orange), 29% (grey)
4	B	106	20% (red), 72% (green), 6% (yellow), 0% (orange), 23% (grey)
4	F	106	17% (red), 73% (green), 5% (yellow), 0% (orange), 23% (grey)
5	C	133	37% (red), 74% (green), 8% (yellow), 0% (orange), 17% (grey)
5	G	133	35% (red), 74% (green), 8% (yellow), 0% (orange), 19% (grey)

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
6	D	129	 <p>24% 63% 12% 25%</p>
6	H	129	 <p>21% 70% 5% 25%</p>

2 Entry composition i

There are 6 unique types of molecules in this entry. The entry contains 23009 atoms, of which 10171 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called Human alpha-satellite DNA (145-MER).

Mol	Chain	Residues	Atoms						AltConf	Trace
			Total	C	H	N	O	P		
1	I	145	5163	1596	1826	603	976	162	18	0

- Molecule 2 is a DNA chain called Human alpha-satellite DNA (145-MER) with a 6-4PP at positions 95-96.

Mol	Chain	Residues	Atoms						AltConf	Trace
			Total	C	H	N	O	P		
2	J	145	5166	1597	1828	605	974	162	18	0

- Molecule 3 is a protein called Histone H3.1.

Mol	Chain	Residues	Atoms						AltConf	Trace
			Total	C	H	N	O	S		
3	A	99	1672	514	857	158	139	4	0	0
3	E	99	1672	514	857	158	139	4	0	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	GLY	-	expression tag	UNP P68431
A	-1	SER	-	expression tag	UNP P68431
A	0	HIS	-	expression tag	UNP P68431
E	-2	GLY	-	expression tag	UNP P68431
E	-1	SER	-	expression tag	UNP P68431
E	0	HIS	-	expression tag	UNP P68431

- Molecule 4 is a protein called Histone H4.

Mol	Chain	Residues	Atoms						AltConf	Trace
			Total	C	H	N	O	S		
4	B	82	1344	412	691	127	113	1	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					AltConf	Trace	
			Total	C	H	N	O			S
4	F	82	1350	412	697	127	113	1	0	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-2	GLY	-	expression tag	UNP P62805
B	-1	SER	-	expression tag	UNP P62805
B	0	HIS	-	expression tag	UNP P62805
F	-2	GLY	-	expression tag	UNP P62805
F	-1	SER	-	expression tag	UNP P62805
F	0	HIS	-	expression tag	UNP P62805

- Molecule 5 is a protein called Histone H2A type 1-B/E.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	H	N	O		
5	C	111	1784	541	924	169	150	0	0
5	G	108	1733	526	898	165	144	0	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	-2	GLY	-	expression tag	UNP P04908
C	-1	SER	-	expression tag	UNP P04908
C	0	HIS	-	expression tag	UNP P04908
G	-2	GLY	-	expression tag	UNP P04908
G	-1	SER	-	expression tag	UNP P04908
G	0	HIS	-	expression tag	UNP P04908

- Molecule 6 is a protein called Histone H2B type 1-J.

Mol	Chain	Residues	Atoms					AltConf	Trace	
			Total	C	H	N	O			S
6	D	97	1562	480	796	142	142	2	0	0
6	H	97	1563	480	797	142	142	2	0	0


There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	-2	GLY	-	expression tag	UNP P06899
D	-1	SER	-	expression tag	UNP P06899
D	0	HIS	-	expression tag	UNP P06899
H	-2	GLY	-	expression tag	UNP P06899
H	-1	SER	-	expression tag	UNP P06899
H	0	HIS	-	expression tag	UNP P06899


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Human alpha-satellite DNA (145-MER)




- Molecule 2: Human alpha-satellite DNA (145-MER) with a 6-4PP at positions 95-96




- Molecule 3: Histone H3.1




- Molecule 3: Histone H3.1




• Molecule 4: Histone H4




• Molecule 4: Histone H4




• Molecule 5: Histone H2A type 1-B/E




• Molecule 5: Histone H2A type 1-B/E





• Molecule 6: Histone H2B type 1-J





• Molecule 6: Histone H2B type 1-J



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	98378	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	40.0	Depositor
Minimum defocus (nm)	Not provided	
Maximum defocus (nm)	Not provided	
Magnification	130000	Depositor
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor
Maximum map value	13.968	Depositor
Minimum map value	-9.715	Depositor
Average map value	0.001	Depositor
Map value standard deviation	0.442	Depositor
Recommended contour level	3.5	Depositor
Map size (Å)	258.0, 258.0, 258.0	wwPDB
Map dimensions	300, 300, 300	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	0.86, 0.86, 0.86	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: T64

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	I	0.94	4/3697 (0.1%)	1.20	8/5699 (0.1%)
2	J	0.94	2/3697 (0.1%)	1.32	12/5694 (0.2%)
3	A	0.52	0/827	0.85	2/1109 (0.2%)
3	E	0.55	0/827	0.81	1/1109 (0.1%)
4	B	0.53	0/660	0.80	0/883
4	F	0.57	0/660	0.81	0/883
5	C	0.48	0/870	0.82	2/1172 (0.2%)
5	G	0.49	0/845	0.76	0/1139
6	D	0.54	0/777	0.73	1/1040 (0.1%)
6	H	0.54	0/777	0.68	0/1040
All	All	0.78	6/13637 (0.0%)	1.08	26/19768 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
3	A	0	1
3	E	0	1
4	B	0	1
4	F	0	1
5	C	0	1
5	G	0	1
6	D	0	1
6	H	0	1
All	All	0	8

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	J	50[B]	DA	O3'-P	-19.92	1.37	1.61
1	I	50[B]	DC	O3'-P	17.27	1.81	1.61
1	I	55[A]	DA	O3'-P	-6.20	1.53	1.61
1	I	55[B]	DA	O3'-P	-6.20	1.53	1.61
2	J	38	DT	C1'-N1	6.13	1.57	1.49

The worst 5 of 26 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	J	95[B]	DT	P-O3'-C3'	39.68	167.32	119.70
1	I	99[A]	DG	P-O3'-C3'	25.65	150.47	119.70
1	I	99[B]	DG	P-O3'-C3'	25.65	150.47	119.70
2	J	99[A]	DG	P-O3'-C3'	24.10	148.62	119.70
2	J	99[B]	DG	P-O3'-C3'	24.10	148.62	119.70

There are no chirality outliers.

5 of 8 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
3	A	135	ARG	Mainchain
4	B	102	GLY	Mainchain
5	C	117	LEU	Peptide
6	D	125	ALA	Mainchain
3	E	135	ARG	Mainchain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	I	3337	1826	1841	10	0
2	J	3338	1828	1841	10	0
3	A	815	857	856	8	0
3	E	815	857	856	5	0
4	B	653	691	696	3	0
4	F	653	697	696	3	0
5	C	860	924	923	7	0
5	G	835	898	897	7	0
6	D	766	796	797	10	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	H	766	797	797	4	0
All	All	12838	10171	10200	45	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 45 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:I:39:DG:O5'	6:D:87:ARG:NH1	2.13	0.82
1:I:52[B]:DC:H42	2:J:94[B]:DA:H61	1.36	0.70
3:E:104:LEU:HD13	3:E:132:ARG:NH2	2.12	0.65
6:H:94:GLU:OE1	6:H:94:GLU:N	2.32	0.62
1:I:119:DT:OP1	5:G:14:LYS:NZ	2.18	0.62

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	A	97/139 (70%)	96 (99%)	1 (1%)	0	100	100
3	E	97/139 (70%)	96 (99%)	1 (1%)	0	100	100
4	B	80/106 (76%)	77 (96%)	3 (4%)	0	100	100
4	F	80/106 (76%)	78 (98%)	2 (2%)	0	100	100
5	C	109/133 (82%)	106 (97%)	2 (2%)	1 (1%)	17	55
5	G	106/133 (80%)	103 (97%)	2 (2%)	1 (1%)	17	55
6	D	95/129 (74%)	89 (94%)	5 (5%)	1 (1%)	14	51
6	H	95/129 (74%)	91 (96%)	3 (3%)	1 (1%)	14	51
All	All	759/1014 (75%)	736 (97%)	19 (2%)	4 (0%)	32	67

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
6	D	105	GLY
6	H	105	GLY
5	C	110	PRO
5	G	110	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	A	86/113 (76%)	84 (98%)	2 (2%)	50	70
3	E	86/113 (76%)	84 (98%)	2 (2%)	50	70
4	B	67/81 (83%)	67 (100%)	0	100	100
4	F	67/81 (83%)	67 (100%)	0	100	100
5	C	88/102 (86%)	88 (100%)	0	100	100
5	G	85/102 (83%)	85 (100%)	0	100	100
6	D	83/107 (78%)	82 (99%)	1 (1%)	71	84
6	H	83/107 (78%)	83 (100%)	0	100	100
All	All	645/806 (80%)	640 (99%)	5 (1%)	82	89

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	A	82	ASP
3	A	130	ARG
6	D	31	LYS
3	E	38	LYS
3	E	130	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
4	F	26	ASN
6	H	96	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	T64	J	95[A]	2	36,44,45	3.31	14 (38%)	47,69,72	4.56	17 (36%)
1	T64	I	95[B]	1	36,44,45	3.35	15 (41%)	47,69,72	5.34	19 (40%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	T64	J	95[A]	2	-	7/22/76/77	0/3/5/5
1	T64	I	95[B]	1	-	7/22/76/77	0/3/5/5

The worst 5 of 29 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	J	95[A]	T64	C2-N3	8.70	1.53	1.38
1	I	95[B]	T64	C2-N3	8.32	1.52	1.38
2	J	95[A]	T64	C2-N1	7.41	1.51	1.36
1	I	95[B]	T64	C2-N1	7.39	1.51	1.36
1	I	95[B]	T64	C4T-N3T	6.80	1.43	1.34

The worst 5 of 36 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	I	95[B]	T64	C4T-C6-N1	-22.21	86.15	110.23
2	J	95[A]	T64	O4T-C5-C4	-15.69	85.51	109.87
1	I	95[B]	T64	O4T-C5-C4	-14.64	87.14	109.87
2	J	95[A]	T64	C4T-C6-N1	-14.01	95.04	110.23
1	I	95[B]	T64	O4T-C5-C5A	-13.77	82.32	109.09

There are no chirality outliers.

5 of 14 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	J	95[A]	T64	O4R-C1R-N1T-C6T
2	J	95[A]	T64	C3'-C4'-C5R-O5R
2	J	95[A]	T64	O4R-C4'-C5R-O5R
2	J	95[A]	T64	C3R-C4R-C5'-O5'
1	I	95[B]	T64	O4R-C1R-N1T-C6T

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-4767. These allow visual inspection of the internal detail of the map and identification of artifacts.

No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map




The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 163



Y Index: 193




Z Index: 156


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal surface views [i](#)


6.4.1 Primary map



X



Y



Z

The images above show the 3D surface view of the map at the recommended contour level 3.5. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


6.5 Mask visualisation

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 68 nm³; this corresponds to an approximate mass of 61 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum i



*Reported resolution corresponds to spatial frequency of 0.250 Å⁻¹


8 Fourier-Shell correlation

This section was not generated. No FSC curve or half-maps provided.

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-4767 and PDB model 6R93. Per-residue inclusion information can be found in section 3 on page 7.

9.1 Map-model overlay [i](#)



The images above show the 3D surface view of the map at the recommended contour level 3.5 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Atom inclusion [i](#)



At the recommended contour level, 68% of all backbone atoms, 65% of all non-hydrogen atoms, are inside the map.