



Full wwPDB X-ray Structure Validation Report ⓘ

Dec 10, 2022 – 01:45 PM EST

PDB ID : 1A2F
Title : PROBING THE STRENGTH AND CHARACTER OF AN ASP-HIS-X HYDROGEN BOND BY INTRODUCING BURIED CHARGES
Authors : Cao, Y.; Goodin, D.B.; Mcree, D.E.
Deposited on : 1998-01-02
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.31.2
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.31.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	A	291	

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 3016 atoms, of which 515 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CYTOCHROME C PEROXIDASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
1	A	291	2866	1502	515	393	451	5	0	0	0

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	53	ILE	THR	engineered mutation	UNP P00431
A	152	GLY	ASP	engineered mutation	UNP P00431
A	172	LYS	MET	engineered mutation	UNP P00431

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	Fe	N	O		
2	A	1	43	34	1	4	4	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	107	Total 107	O 107	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: CYTOCHROME C PEROXIDASE



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	103.80Å 73.50Å 44.90Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	5.00 – 2.10 14.92 – 1.91	Depositor EDS
% Data completeness (in resolution range)	(Not available) (5.00-2.10) 78.9 (14.92-1.91)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	0.09	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.58 (at 1.91Å)	Xtrriage
Refinement program	XTALVIEW, X-PLOR 3.1	Depositor
R, R_{free}	0.184 , (Not available) 0.484 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	20.2	Xtrriage
Anisotropy	0.644	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 47.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.62	EDS
Total number of atoms	3016	wwPDB-VP
Average B, all atoms (Å ²)	18.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.28% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: HEM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.86	0/2417	1.58	40/3271 (1.2%)

There are no bond length outliers.

All (40) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	31	ARG	NE-CZ-NH2	-12.81	113.89	120.30
1	A	51	TRP	CD1-CG-CD2	9.44	113.85	106.30
1	A	31	ARG	NE-CZ-NH1	8.96	124.78	120.30
1	A	51	TRP	CE2-CD2-CG	-8.24	100.71	107.30
1	A	191	TRP	CE2-CD2-CG	-8.17	100.77	107.30
1	A	101	TRP	CD1-CG-CD2	8.02	112.72	106.30
1	A	143	ARG	NE-CZ-NH1	7.75	124.18	120.30
1	A	191	TRP	CD1-CG-CD2	7.70	112.46	106.30
1	A	211	TRP	CD1-CG-CD2	7.46	112.27	106.30
1	A	223	TRP	CD1-CG-CD2	7.46	112.27	106.30
1	A	126	TRP	CD1-CG-CD2	7.27	112.11	106.30
1	A	51	TRP	CB-CG-CD1	-7.24	117.59	127.00
1	A	101	TRP	CE2-CD2-CG	-7.17	101.56	107.30
1	A	127	ARG	NE-CZ-NH2	-7.10	116.75	120.30
1	A	57	TRP	CD1-CG-CD2	6.83	111.76	106.30
1	A	223	TRP	CE2-CD2-CG	-6.81	101.85	107.30
1	A	48	ARG	NE-CZ-NH1	6.67	123.64	120.30
1	A	51	TRP	CG-CD2-CE3	6.62	139.86	133.90
1	A	211	TRP	CE2-CD2-CG	-6.60	102.02	107.30
1	A	51	TRP	CG-CD1-NE1	-6.52	103.58	110.10
1	A	229	TYR	CB-CG-CD2	-6.47	117.12	121.00
1	A	57	TRP	CE2-CD2-CG	-6.46	102.13	107.30
1	A	127	ARG	NE-CZ-NH1	6.33	123.47	120.30
1	A	126	TRP	CE2-CD2-CG	-6.24	102.31	107.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	278	LYS	CA-CB-CG	-6.16	99.84	113.40
1	A	162	ASN	CA-C-N	-5.99	104.03	117.20
1	A	251	TYR	CB-CG-CD2	-5.84	117.49	121.00
1	A	101	TRP	CG-CD1-NE1	-5.78	104.32	110.10
1	A	187	TYR	CB-CG-CD2	-5.77	117.54	121.00
1	A	155	ARG	NE-CZ-NH2	-5.71	117.44	120.30
1	A	230	MET	CA-CB-CG	5.70	122.99	113.30
1	A	163	MET	CG-SD-CE	-5.55	91.31	100.20
1	A	71	TYR	CB-CG-CD2	-5.49	117.70	121.00
1	A	277	PRO	CA-C-N	5.42	129.12	117.20
1	A	166	ARG	NE-CZ-NH2	-5.25	117.68	120.30
1	A	191	TRP	CB-CG-CD1	-5.24	120.19	127.00
1	A	39	TYR	CB-CG-CD2	-5.22	117.87	121.00
1	A	101	TRP	CG-CD2-CE3	5.16	138.55	133.90
1	A	230	MET	CG-SD-CE	-5.09	92.06	100.20
1	A	101	TRP	CB-CG-CD1	-5.08	120.39	127.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2351	515	2232	18	23
2	A	43	0	30	0	0
3	A	107	0	0	2	13
All	All	2501	515	2262	18	23

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (18) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:168:VAL:O	1:A:172:LYS:HG2	1.98	0.63

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:195:ASN:HD22	1:A:195:ASN:H	1.47	0.63
1:A:206:LEU:HD13	1:A:231:MET:SD	2.38	0.62
1:A:20:GLN:HE22	1:A:287:LYS:H	1.51	0.58
1:A:74:LYS:HD2	1:A:74:LYS:H	1.70	0.57
1:A:200:ASN:H	1:A:255:GLN:HE21	1.56	0.53
1:A:130:ARG:NE	3:A:326:HOH:O	2.43	0.51
1:A:256:ASP:O	1:A:260:LYS:HD2	2.17	0.44
1:A:164:ASN:O	1:A:168:VAL:HG23	2.17	0.44
1:A:20:GLN:HE22	1:A:287:LYS:N	2.13	0.44
1:A:218:ALA:O	1:A:219:ASN:HB2	2.18	0.44
1:A:195:ASN:H	1:A:195:ASN:ND2	2.14	0.43
1:A:125:PRO:HG3	1:A:285:ILE:HD11	2.01	0.42
1:A:113:VAL:O	1:A:117:GLN:HG3	2.20	0.41
1:A:195:ASN:HD22	1:A:195:ASN:N	2.18	0.41
1:A:4:LEU:HB2	1:A:62:ASN:HB3	2.02	0.41
1:A:130:ARG:CZ	3:A:326:HOH:O	2.69	0.40
1:A:166:ARG:HH21	1:A:257:LYS:HZ2	1.68	0.40

All (23) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:14:ARG:HH21	1:A:213:LEU:O[4_5610]	1.01	0.59
1:A:14:ARG:HH21	1:A:213:LEU:C[4_5610]	1.01	0.59
1:A:212:LYS:NZ	3:A:374:HOH:O[4_4610]	1.70	0.50
1:A:212:LYS:CG	3:A:372:HOH:O[4_4610]	1.70	0.50
1:A:212:LYS:CD	3:A:372:HOH:O[4_4610]	1.74	0.46
1:A:14:ARG:NH2	1:A:213:LEU:C[4_5610]	1.81	0.39
1:A:264:LYS:HZ1	3:A:389:HOH:O[3_559]	1.22	0.38
1:A:264:LYS:NZ	3:A:389:HOH:O[3_559]	1.83	0.37
1:A:14:ARG:NH2	1:A:213:LEU:O[4_5610]	1.86	0.34
1:A:212:LYS:CE	3:A:374:HOH:O[4_4610]	1.86	0.34
1:A:12:LYS:CB	1:A:214:GLU:OE2[4_5610]	1.87	0.33
1:A:214:GLU:OE2	3:A:374:HOH:O[4_4610]	1.89	0.31
1:A:212:LYS:HZ3	3:A:374:HOH:O[4_4610]	1.35	0.25
1:A:212:LYS:CE	3:A:372:HOH:O[4_4610]	1.97	0.23
1:A:14:ARG:NH2	1:A:214:GLU:N[4_5610]	2.02	0.18
1:A:214:GLU:CG	3:A:373:HOH:O[4_4610]	2.05	0.15
1:A:101:TRP:CB	1:A:213:LEU:O[4_5610]	2.07	0.13
1:A:14:ARG:HH21	1:A:214:GLU:N[4_5610]	1.54	0.06

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:12:LYS:N	1:A:214:GLU:OE2[4_5610]	2.15	0.05
1:A:12:LYS:H	1:A:214:GLU:OE2[4_5610]	1.56	0.04
1:A:214:GLU:CB	3:A:373:HOH:O[4_4610]	2.17	0.03
1:A:213:LEU:O	3:A:373:HOH:O[4_4610]	2.18	0.02
1:A:293:GLY:O	3:A:330:HOH:O[3_549]	2.18	0.02

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	289/291 (99%)	282 (98%)	6 (2%)	1 (0%)	41 41

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	162	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	249/249 (100%)	240 (96%)	9 (4%)	35 36

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	28	LEU
1	A	35	GLU
1	A	74	LYS
1	A	80	PRO
1	A	159	GLN
1	A	166	ARG
1	A	171	LEU
1	A	195	ASN
1	A	260	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	20	GLN
1	A	24	ASN
1	A	87	ASN
1	A	195	ASN
1	A	208	ASN
1	A	220	ASN
1	A	240	GLN
1	A	255	GLN
1	A	292	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	HEM	A	1	3,1	41,50,50	1.95	16 (39%)	45,82,82	1.62	4 (8%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	A	1	3,1	-	4/12/54/54	-

All (16) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1	HEM	C4D-ND	-4.16	1.33	1.40
2	A	1	HEM	C4D-C3D	-3.84	1.38	1.45
2	A	1	HEM	C1D-C2D	-3.32	1.38	1.44
2	A	1	HEM	C1B-NB	-3.28	1.34	1.40
2	A	1	HEM	C3C-CAC	-3.17	1.41	1.47
2	A	1	HEM	CAB-C3B	-3.00	1.39	1.47
2	A	1	HEM	C3B-C4B	-2.89	1.39	1.44
2	A	1	HEM	C1B-C2B	-2.76	1.39	1.44
2	A	1	HEM	C1D-ND	-2.71	1.33	1.38
2	A	1	HEM	CBC-CAC	2.62	1.46	1.29
2	A	1	HEM	CBB-CAB	2.60	1.43	1.30
2	A	1	HEM	C4B-NB	-2.57	1.33	1.38
2	A	1	HEM	FE-ND	2.45	2.09	1.96
2	A	1	HEM	CHB-C1B	2.21	1.40	1.35
2	A	1	HEM	O2A-CGA	-2.12	1.23	1.30
2	A	1	HEM	O2D-CGD	-2.10	1.23	1.30

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1	HEM	C4B-CHC-C1C	5.73	130.12	122.56

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1	HEM	C4C-CHD-C1D	4.88	128.99	122.56
2	A	1	HEM	CMC-C2C-C3C	2.91	130.12	124.68
2	A	1	HEM	O2A-CGA-CBA	2.15	120.94	114.03

There are no chirality outliers.


All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	1	HEM	CAA-CBA-CGA-O2A
2	A	1	HEM	CAA-CBA-CGA-O1A
2	A	1	HEM	CAD-CBD-CGD-O2D
2	A	1	HEM	CAD-CBD-CGD-O1D

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.


6.3 Carbohydrates

Unable to reproduce the depositors R factor - this section is therefore empty.

6.4 Ligands

Unable to reproduce the depositors R factor - this section is therefore empty.

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.