



# Full wwPDB X-ray Structure Validation Report ⓘ

Aug 16, 2023 – 01:35 AM EDT

PDB ID : 1YBO  
Title : Crystal structure of the PDZ tandem of human syntenin with syndecan peptide  
Authors : Grembecka, J.; Cooper, D.R.; Cierpicki, T.; Kang, B.S.; Devedjiev, Y.; Derewenda, Z.  
Deposited on : 2004-12-21  
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.35  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	166	
1	B	166	
2	C	17	
2	D	17	

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 2770 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Syntenin 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	163	1244	778	222	238	6	0	0	0
1	B	164	1245	777	223	238	7	0	0	0

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	108	GLY	-	cloning artifact	UNP O00560
A	109	ALA	-	cloning artifact	UNP O00560
A	110	MET	-	cloning artifact	UNP O00560
A	111	ASP	-	cloning artifact	UNP O00560
A	112	PRO	-	cloning artifact	UNP O00560
B	108	GLY	-	cloning artifact	UNP O00560
B	109	ALA	-	cloning artifact	UNP O00560
B	110	MET	-	cloning artifact	UNP O00560
B	111	ASP	-	cloning artifact	UNP O00560
B	112	PRO	-	cloning artifact	UNP O00560

- Molecule 2 is a protein called Syndecan-4.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	C	6	53	34	7	12	0	0	0
2	D	4	38	26	4	8	0	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	92	Total	O	0	0
			92	92		

*Continued on next page...*


*Continued from previous page...*


<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
3	B	90	Total	O	0	0
			90	90		
3	C	4	Total	O	0	0
			4	4		
3	D	4	Total	O	0	0
			4	4		

### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Syntenin 1

Chain A: 




- Molecule 1: Syntenin 1

Chain B: 




- Molecule 2: Syndecan-4

Chain C: 



- Molecule 2: Syndecan-4

Chain D: 



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	72.32Å 72.32Å 125.92Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.00 – 2.30 36.30 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.3 (40.00-2.30) 99.3 (36.30-2.30)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	13.74 (at 2.29Å)	Xtrriage
Refinement program	REFMAC	Depositor
R, $R_{free}$	0.193 , 0.273 0.207 , 0.291	Depositor DCC
$R_{free}$ test set	800 reflections (5.20%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	27.2	Xtrriage
Anisotropy	0.058	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 49.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.44$ , $\langle L^2 \rangle = 0.27$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	2770	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	28.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.96% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.53	0/1260	1.41	13/1696 (0.8%)
1	B	0.54	0/1260	1.33	11/1697 (0.6%)
2	C	0.72	0/54	1.33	0/71
2	D	0.56	0/39	1.13	0/50
All	All	0.54	0/2613	1.36	24/3514 (0.7%)

There are no bond length outliers.

All (24) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	193	ARG	NE-CZ-NH2	-10.44	115.08	120.30
1	A	256	ASP	CB-CG-OD1	-9.15	110.07	118.30
1	A	193	ARG	NE-CZ-NH1	8.43	124.52	120.30
1	B	197	ARG	NE-CZ-NH1	8.03	124.31	120.30
1	B	204	ASP	CB-CG-OD1	7.67	125.20	118.30
1	B	197	ARG	NE-CZ-NH2	-7.61	116.50	120.30
1	A	256	ASP	CB-CG-OD2	7.39	124.95	118.30
1	B	128	ARG	NE-CZ-NH1	7.34	123.97	120.30
1	B	156	ASP	CB-CG-OD2	-7.24	111.79	118.30
1	A	229	ARG	NE-CZ-NH1	6.66	123.63	120.30
1	B	133	ASP	CB-CG-OD1	6.66	124.29	118.30
1	A	261	SER	C-N-CA	-6.47	108.70	122.30
1	B	122	ASP	CB-CG-OD1	6.45	124.11	118.30
1	A	172	ASP	CB-CG-OD1	6.36	124.03	118.30
1	B	156	ASP	CB-CG-OD1	6.24	123.92	118.30
1	B	113	ARG	NE-CZ-NH1	5.99	123.30	120.30
1	A	270	MET	CG-SD-CE	-5.85	90.84	100.20
1	B	113	ARG	NE-CZ-NH2	-5.77	117.42	120.30
1	A	153	ARG	NE-CZ-NH2	-5.63	117.48	120.30
1	A	128	ARG	NE-CZ-NH2	-5.56	117.52	120.30
1	A	204	ASP	CB-CG-OD1	5.44	123.19	118.30
1	A	133	ASP	CB-CG-OD2	-5.34	113.50	118.30
1	B	210	GLY	N-CA-C	5.16	126.00	113.10

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	229	ARG	CD-NE-CZ	5.03	130.64	123.60

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1244	0	1271	7	0
1	B	1245	0	1276	10	0
2	C	53	0	44	2	0
2	D	38	0	28	0	0
3	A	92	0	0	5	1
3	B	90	0	0	3	0
3	C	4	0	0	0	0
3	D	4	0	0	0	0
All	All	2770	0	2619	19	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (19) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:268:THR:HB	3:B:351:HOH:O	1.84	0.77
2:C:1:THR:HB	2:C:3:GLU:H	1.67	0.58
1:B:268:THR:CB	3:B:351:HOH:O	2.44	0.57
1:A:181:ALA:HA	3:A:298:HOH:O	2.06	0.56
1:B:196:GLU:CG	1:B:268:THR:CG2	2.89	0.50
1:A:210:GLY:HA3	1:A:225:SER:HB2	1.95	0.48
1:A:193:ARG:HD3	1:A:270:MET:CE	2.43	0.48
1:B:196:GLU:HG2	1:B:268:THR:CG2	2.44	0.47
1:B:210:GLY:HA3	1:B:225:SER:HB2	1.98	0.45
1:A:270:MET:HA	3:A:359:HOH:O	2.17	0.44
1:B:115:VAL:HG23	1:B:117:LEU:HD11	1.98	0.44

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:261:SER:O	3:A:280:HOH:O	2.20	0.44
1:B:176:LYS:HE3	1:B:180:GLN:NE2	2.33	0.43
1:B:206:THR:O	3:B:333:HOH:O	2.21	0.43
1:A:203:LYS:HG3	3:A:352:HOH:O	2.19	0.42
1:A:120:ASP:HB2	3:A:278:HOH:O	2.20	0.41
1:B:196:GLU:CD	1:B:268:THR:HG21	2.42	0.40
1:B:176:LYS:NZ	1:B:180:GLN:NE2	2.69	0.40
2:C:1:THR:HB	2:C:3:GLU:N	2.33	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:282:HOH:O	3:A:350:HOH:O[3_645]	2.16	0.04

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	161/166 (97%)	153 (95%)	5 (3%)	3 (2%)	8	7
1	B	162/166 (98%)	159 (98%)	2 (1%)	1 (1%)	25	31
2	C	4/17 (24%)	3 (75%)	1 (25%)	0	100	100
2	D	2/17 (12%)	2 (100%)	0	0	100	100
All	All	329/366 (90%)	317 (96%)	8 (2%)	4 (1%)	13	14

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	124	LYS
1	A	122	ASP

Continued on next page...

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	193	ARG
1	B	193	ARG

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	139/140 (99%)	135 (97%)	4 (3%)	42	58
1	B	139/140 (99%)	137 (99%)	2 (1%)	67	81
2	C	5/14 (36%)	4 (80%)	1 (20%)	1	1
2	D	3/14 (21%)	3 (100%)	0	100	100
All	All	286/308 (93%)	279 (98%)	7 (2%)	49	66

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	118	CYS
1	A	179	LYS
1	A	187	THR
1	A	267	ILE
1	B	235	GLU
1	B	252	SER
2	C	1	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	180	GLN
1	B	237	ASN
1	B	253	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	163/166 (98%)	0.00	4 (2%) 57 64	19, 26, 40, 48	0
1	B	164/166 (98%)	-0.02	5 (3%) 50 57	19, 27, 36, 50	2 (1%)
2	C	6/17 (35%)	1.23	1 (16%) 1 2	25, 31, 40, 42	0
2	D	4/17 (23%)	0.43	0 100 100	35, 36, 37, 40	0
All	All	337/366 (92%)	0.02	10 (2%) 50 57	19, 27, 40, 50	2 (0%)

All (10) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	109	ALA	6.8
2	C	1	THR	5.2
1	B	110	MET	3.7
1	A	124	LYS	3.7
1	A	121	GLN	3.6
1	A	181	ALA	2.9
1	B	248	GLY	2.9
1	B	184	GLU	2.8
1	B	205	SER	2.3
1	A	122	ASP	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.