



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 12, 2024 – 10:33 PM EDT

PDB ID : 4C50  
Title : Crystal Structure of the Catalase-Peroxidase (KatG) D137S mutant from Mycobacterium Tuberculosis  
Authors : Hersleth, H.-P.; Zhao, X.; Magliozzo, R.S.; Andersson, K.K.  
Deposited on : 2013-09-10  
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtrriage (Phenix) : 1.20.1  
EDS : 2.36.2  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4661 (2.50-2.50)
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	740	 2% 87% 9%
1	B	740	 11% 89% 8%

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 11241 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called CATALASE-PEROXIDASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	716	5524	3512	952	1041	19	0	0	0
1	B	716	5533	3517	953	1044	19	0	1	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	137	SER	ASP	engineered mutation	UNP Q08129
B	137	SER	ASP	engineered mutation	UNP Q08129

- Molecule 2 is alpha-D-glucopyranose (three-letter code: GLC) (formula: C<sub>6</sub>H<sub>12</sub>O<sub>6</sub>).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
2	A	1	12	6	6	0	0

- Molecule 3 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula:  $C_{34}H_{32}FeN_4O_4$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
3	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
3	B	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

- Molecule 4 is ACETATE ION (three-letter code: ACT) (formula:  $C_2H_3O_2$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	2	2		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	56	Total	O	0	0
			56	56		
5	B	26	Total	O	0	0
			26	26		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: CATALASE-PEROXIDASE



- Molecule 1: CATALASE-PEROXIDASE



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 42 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	150.59Å 150.59Å 157.35Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	75.41 – 2.50 69.73 – 2.50	Depositor EDS
% Data completeness (in resolution range)	99.8 (75.41-2.50) 99.8 (69.73-2.50)	Depositor EDS
$R_{merge}$	0.13	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.71 (at 2.51Å)	Xtriage
Refinement program	REFMAC 5.7.0032	Depositor
R, $R_{free}$	0.264 , 0.311 0.267 , 0.312	Depositor DCC
$R_{free}$ test set	3188 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	27.4	Xtriage
Anisotropy	0.047	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 29.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.019 for -h,l,k 0.013 for -l,-k,-h	Xtriage
$F_o, F_c$ correlation	0.87	EDS
Total number of atoms	11241	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.29% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: HEM, ACT, GLC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.61	0/5675	0.71	1/7724 (0.0%)
1	B	0.55	0/5684	0.70	1/7736 (0.0%)
All	All	0.58	0/11359	0.71	2/15460 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	63	ASP	CB-CG-OD2	-5.38	113.46	118.30
1	A	632	ARG	NE-CZ-NH2	-5.25	117.68	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5524	0	5353	38	0
1	B	5533	0	5358	29	0
2	A	12	0	12	0	0
3	A	43	0	30	2	0
3	B	43	0	30	1	0
4	A	4	0	3	0	0
5	A	56	0	0	1	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	B	26	0	0	1	0
All	All	11241	0	10786	67	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (67) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:107:TRP:HH2	1:A:229:TYR:CE2	1.15	1.62
1:B:107:TRP:HH2	1:B:229:TYR:CE2	1.15	1.57
1:B:107:TRP:CH2	1:B:229:TYR:HE2	1.23	1.53
1:A:107:TRP:CH2	1:A:229:TYR:CE2	1.99	1.48
1:A:107:TRP:CH2	1:A:229:TYR:HE2	1.28	1.47
1:B:107:TRP:CH2	1:B:229:TYR:CE2	1.96	1.42
1:B:107:TRP:HH2	1:B:229:TYR:CD2	1.83	0.96
1:A:107:TRP:HH2	1:A:229:TYR:CD2	1.86	0.92
1:B:107:TRP:CZ3	1:B:229:TYR:HE2	1.88	0.91
1:A:107:TRP:CZ3	1:A:229:TYR:HE2	1.95	0.85
1:A:258:ASN:C	1:A:258:ASN:HD22	1.86	0.79
1:A:49:HIS:O	1:A:49:HIS:CD2	2.41	0.73
1:B:69:ALA:HB2	5:B:2003:HOH:O	1.88	0.72
1:B:107:TRP:CZ3	1:B:229:TYR:CE2	2.70	0.72
1:A:107:TRP:CZ3	1:A:229:TYR:CE2	2.74	0.70
1:A:49:HIS:O	1:A:49:HIS:HD2	1.75	0.69
1:B:107:TRP:CH2	1:B:229:TYR:CD2	2.68	0.63
1:A:107:TRP:CH2	1:A:229:TYR:CD2	2.71	0.62
1:B:258:ASN:C	1:B:258:ASN:HD22	2.04	0.61
1:A:701:ASN:HD22	1:A:704:LEU:H	1.49	0.61
1:B:45:LEU:HD23	1:B:611:LEU:HD21	1.84	0.59
1:A:318:GLU:H	1:A:352:GLN:HE22	1.51	0.58
1:A:45:LEU:HD23	1:A:611:LEU:HD21	1.87	0.57
1:B:275:THR:HG22	3:B:1741:HEM:HAA1	1.86	0.56
1:B:606:ALA:HB1	1:B:694:VAL:HG23	1.90	0.54
1:B:119:ARG:HH21	1:B:615:ASN:HD22	1.57	0.52
1:A:91:TRP:CE3	1:A:143:LYS:HD2	2.45	0.52
1:B:318:GLU:H	1:B:352:GLN:HE22	1.57	0.51
1:A:139:ALA:HA	1:A:300:TRP:CH2	2.45	0.51
1:A:139:ALA:HA	1:A:300:TRP:CZ3	2.46	0.51
1:A:116:HIS:CD2	1:A:219:PRO:HB3	2.46	0.50
1:B:345:LYS:HE2	1:B:351:TRP:CH2	2.46	0.49

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:91:TRP:CE3	1:B:143:LYS:HD2	2.48	0.49
1:A:119:ARG:HH21	1:A:615:ASN:HD22	1.61	0.49
1:B:45:LEU:CD2	1:B:611:LEU:HD21	2.43	0.48
1:A:258:ASN:C	1:A:258:ASN:ND2	2.59	0.48
1:B:49:HIS:O	1:B:49:HIS:CD2	2.67	0.48
1:A:45:LEU:CD2	1:A:611:LEU:HD21	2.44	0.47
1:A:259:ASP:HA	5:A:2023:HOH:O	2.14	0.47
1:A:107:TRP:CD1	1:A:108:HIS:HD2	2.33	0.47
1:A:606:ALA:HB1	1:A:694:VAL:HG23	1.96	0.46
1:B:95:TYR:CE1	1:B:367:PRO:HA	2.51	0.45
1:A:239:PRO:O	1:A:241:PRO:HD3	2.16	0.45
1:B:107:TRP:CD1	1:B:108:HIS:HD2	2.35	0.45
1:B:701:ASN:HD22	1:B:704:LEU:H	1.64	0.45
1:A:378:LEU:HB2	1:A:381:ASP:OD2	2.17	0.45
1:B:116:HIS:CD2	1:B:219:PRO:HB3	2.52	0.45
1:A:650:SER:O	1:A:651:GLU:C	2.56	0.44
1:A:666:ILE:HA	1:A:681:LYS:O	2.18	0.43
1:A:39:TRP:HB2	1:A:42:ARG:HD2	2.01	0.43
1:A:95:TYR:CE1	1:A:367:PRO:HA	2.53	0.43
1:A:345:LYS:HE2	1:A:351:TRP:CH2	2.53	0.43
1:B:616:LEU:HD23	1:B:616:LEU:HA	1.86	0.42
1:A:100:PRO:HB2	3:A:1741:HEM:O2D	2.18	0.42
1:B:317:ILE:HG23	1:B:352:GLN:HE21	1.84	0.42
1:A:646:PHE:O	1:A:656:ASP:HB3	2.21	0.41
1:A:672:PRO:O	1:A:674:ASP:N	2.53	0.41
1:B:655:ASN:O	1:B:656:ASP:C	2.59	0.41
1:A:275:THR:OG1	1:A:317:ILE:O	2.35	0.41
1:A:314:THR:HB	3:A:1741:HEM:O1A	2.21	0.41
1:A:658:PHE:HD1	1:A:708:VAL:HG13	1.86	0.41
1:B:67:GLU:HB3	1:B:158:LYS:HA	2.03	0.41
1:A:317:ILE:HG23	1:A:352:GLN:HE21	1.85	0.41
1:A:616:LEU:HD23	1:A:616:LEU:HA	1.83	0.41
1:B:95:TYR:CD1	1:B:367:PRO:HA	2.56	0.41
1:B:167:PHE:CE2	1:B:171:CYS:SG	3.14	0.40
1:B:628:VAL:O	1:B:632:ARG:HG2	2.21	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	714/740 (96%)	677 (95%)	32 (4%)	5 (1%)	22	39
1	B	715/740 (97%)	673 (94%)	37 (5%)	5 (1%)	22	39
All	All	1429/1480 (97%)	1350 (94%)	69 (5%)	10 (1%)	22	39

All (10) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	512	GLY
1	A	673	ALA
1	B	512	GLY
1	B	513	ASP
1	B	673	ALA
1	A	513	ASP
1	A	514	LEU
1	B	656	ASP
1	A	370	GLY
1	B	514	LEU

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	565/584 (97%)	546 (97%)	19 (3%)	37	63
1	B	566/584 (97%)	545 (96%)	21 (4%)	34	60
All	All	1131/1168 (97%)	1091 (96%)	40 (4%)	36	62

All (40) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	82	GLU
1	A	175	SER
1	A	179	LYS
1	A	195	GLU
1	A	218	ASN
1	A	258	ASN
1	A	310	LYS
1	A	327	LYS
1	A	373	ARG
1	A	376	THR
1	A	433	LYS
1	A	457	SER
1	A	526	GLU
1	A	554	LYS
1	A	562	ASN
1	A	564	THR
1	A	590	LYS
1	A	600	LYS
1	A	640	ARG
1	B	81[A]	GLU
1	B	81[B]	GLU
1	B	82	GLU
1	B	175	SER
1	B	179	LYS
1	B	195	GLU
1	B	218	ASN
1	B	258	ASN
1	B	310	LYS
1	B	327	LYS
1	B	330	ASN
1	B	373	ARG
1	B	376	THR
1	B	433	LYS
1	B	526	GLU
1	B	554	LYS
1	B	562	ASN
1	B	564	THR
1	B	590	LYS
1	B	600	LYS
1	B	640	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (27)

such sidechains are listed below:

Mol	Chain	Res	Type
1	A	25	HIS
1	A	36	GLN
1	A	49	HIS
1	A	51	ASN
1	A	116	HIS
1	A	218	ASN
1	A	258	ASN
1	A	330	ASN
1	A	352	GLN
1	A	447	HIS
1	A	535	ASN
1	A	615	ASN
1	A	701	ASN
1	B	36	GLN
1	B	49	HIS
1	B	51	ASN
1	B	116	HIS
1	B	218	ASN
1	B	258	ASN
1	B	330	ASN
1	B	352	GLN
1	B	447	HIS
1	B	535	ASN
1	B	602	ASN
1	B	615	ASN
1	B	701	ASN
1	B	717	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	ACT	A	1743	-	3,3,3	0.96	0	3,3,3	0.45	0
3	HEM	A	1741	5,1	42,50,50	2.03	11 (26%)	46,82,82	1.73	9 (19%)
2	GLC	A	1742	-	12,12,12	0.92	0	17,17,17	2.24	7 (41%)
3	HEM	B	1741	1	42,50,50	1.96	8 (19%)	46,82,82	1.84	15 (32%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	HEM	A	1741	5,1	-	2/12/54/54	-
2	GLC	A	1742	-	-	0/2/22/22	0/1/1/1
3	HEM	B	1741	1	-	2/12/54/54	-

All (19) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	1741	HEM	C3D-C2D	7.04	1.51	1.36
3	A	1741	HEM	C3D-C2D	6.50	1.50	1.36
3	B	1741	HEM	C3C-C2C	-5.17	1.33	1.40
3	A	1741	HEM	C3C-C2C	-4.40	1.34	1.40
3	B	1741	HEM	FE-ND	3.86	2.19	1.98
3	A	1741	HEM	FE-ND	3.57	2.18	1.98
3	A	1741	HEM	C3C-C4C	3.52	1.46	1.41
3	A	1741	HEM	CHC-C4B	-3.01	1.32	1.40
3	B	1741	HEM	C3C-CAC	2.86	1.54	1.47
3	A	1741	HEM	CAB-C3B	2.82	1.54	1.47
3	B	1741	HEM	CAB-C3B	2.78	1.54	1.47
3	A	1741	HEM	C1B-NB	-2.76	1.35	1.40

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1741	HEM	C3B-C2B	-2.63	1.31	1.37
3	B	1741	HEM	C3C-C4C	2.60	1.45	1.41
3	B	1741	HEM	CMB-C2B	2.55	1.56	1.50
3	A	1741	HEM	C3C-CAC	2.45	1.53	1.47
3	A	1741	HEM	C1A-NA	2.04	1.40	1.36
3	A	1741	HEM	C2A-C3A	-2.02	1.31	1.37
3	B	1741	HEM	CHC-C4B	-2.00	1.35	1.40

All (31) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	1741	HEM	C4D-ND-C1D	5.22	111.39	105.21
3	A	1741	HEM	CAA-CBA-CGA	-4.45	101.84	113.83
3	A	1741	HEM	CAD-C3D-C4D	4.25	132.11	124.70
2	A	1742	GLC	O2-C2-C1	4.03	118.54	109.25
3	A	1741	HEM	C3B-C4B-NB	3.79	112.19	109.47
2	A	1742	GLC	O3-C3-C2	-3.71	101.62	110.38
2	A	1742	GLC	O6-C6-C5	3.67	123.82	111.33
2	A	1742	GLC	O5-C5-C6	3.62	115.41	106.44
3	B	1741	HEM	C2C-C3C-C4C	3.61	109.42	106.90
3	A	1741	HEM	C4D-ND-C1D	3.43	109.26	105.21
2	A	1742	GLC	O5-C1-C2	3.23	115.97	110.30
3	B	1741	HEM	C4C-CHD-C1D	3.12	126.67	122.56
3	B	1741	HEM	CAD-C3D-C4D	3.07	130.04	124.70
3	B	1741	HEM	CAA-CBA-CGA	-2.95	105.90	113.83
2	A	1742	GLC	C3-C4-C5	2.82	115.35	110.23
3	B	1741	HEM	CMA-C3A-C4A	-2.81	124.33	128.46
3	A	1741	HEM	CMD-C2D-C1D	2.64	129.17	125.03
3	B	1741	HEM	C3B-C4B-NB	2.64	111.37	109.47
3	A	1741	HEM	CMC-C2C-C3C	2.48	129.64	124.68
3	B	1741	HEM	C4A-C3A-C2A	2.44	108.70	107.00
3	B	1741	HEM	C4B-CHC-C1C	-2.37	119.44	122.56
3	A	1741	HEM	C4D-C3D-C2D	-2.29	103.56	106.89
2	A	1742	GLC	O3-C3-C4	2.28	115.74	110.38
3	B	1741	HEM	CHD-C1D-ND	2.25	126.86	124.44
3	A	1741	HEM	CHB-C1B-NB	-2.22	121.62	124.37
3	B	1741	HEM	O1D-CGD-CBD	-2.20	116.12	123.09
3	B	1741	HEM	CHC-C4B-NB	-2.18	122.09	124.44
3	B	1741	HEM	C4D-C3D-C2D	-2.11	103.81	106.89
3	B	1741	HEM	O2D-CGD-CBD	2.10	120.63	114.00
3	B	1741	HEM	CMC-C2C-C3C	2.06	128.79	124.68
3	A	1741	HEM	C4A-C3A-C2A	2.04	108.42	107.00

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	1741	HEM	CAA-CBA-CGA-O2A
3	A	1741	HEM	CAA-CBA-CGA-O1A
3	B	1741	HEM	C4B-C3B-CAB-CBB
3	B	1741	HEM	C2B-C3B-CAB-CBB


There are no ring outliers.


2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1741	HEM	2	0
3	B	1741	HEM	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	716/740 (96%)	0.10	18 (2%) 57 61	9, 25, 48, 82	0
1	B	716/740 (96%)	0.80	84 (11%) 4 4	13, 39, 75, 129	0
All	All	1432/1480 (96%)	0.45	102 (7%) 16 16	9, 31, 67, 129	0

All (102) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	532	ALA	7.8
1	B	533	PRO	5.6
1	B	683	GLY	5.1
1	B	535	ASN	5.0
1	A	510	PRO	4.9
1	B	514	LEU	4.7
1	B	541	ALA	4.6
1	B	534	GLY	4.5
1	B	581	VAL	4.5
1	B	517	VAL	4.1
1	B	462	ILE	3.9
1	B	561	HIS	3.9
1	B	455	ILE	3.7
1	B	452	GLU	3.7
1	B	467	LEU	3.6
1	B	497	ILE	3.4
1	B	459	LYS	3.4
1	B	531	ALA	3.4
1	A	369	GLY	3.3
1	B	550	ALA	3.2
1	B	469	VAL	3.1
1	A	673	ALA	3.1
1	B	536	ILE	3.1
1	B	530	SER	3.1

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
1	A	297	GLY	3.1
1	B	503	VAL	3.1
1	A	511	ASP	3.0
1	B	510	PRO	3.0
1	B	650	SER	3.0
1	B	463	ARG	3.0
1	A	97	HIS	3.0
1	B	498	ARG	3.0
1	B	494	GLY	2.9
1	B	526	GLU	2.9
1	B	423	VAL	2.9
1	B	473	VAL	2.9
1	B	474	SER	2.9
1	B	472	LEU	2.9
1	B	475	THR	2.9
1	B	527	SER	2.8
1	B	554	LYS	2.8
1	A	96	GLY	2.8
1	B	511	ASP	2.8
1	B	555	ALA	2.8
1	B	564	THR	2.8
1	B	216	LEU	2.8
1	B	524	ILE	2.8
1	B	528	PHE	2.7
1	A	147	LEU	2.7
1	B	538	VAL	2.7
1	B	508	ASN	2.7
1	B	673	ALA	2.7
1	A	310	LYS	2.7
1	B	357	ASP	2.7
1	B	476	ALA	2.6
1	B	654	THR	2.6
1	B	212	GLY	2.6
1	B	521	LEU	2.6
1	B	629	GLY	2.6
1	B	678	TYR	2.5
1	B	445	VAL	2.5
1	B	499	LEU	2.5
1	B	738	ASP	2.5
1	B	447	HIS	2.5
1	A	533	PRO	2.5
1	B	525	GLN	2.5

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	684	SER	2.4
1	B	519	ARG	2.4
1	B	633	VAL	2.3
1	A	399	GLU	2.3
1	B	648	GLU	2.3
1	B	563	ILE	2.3
1	B	206	GLY	2.3
1	B	540	PHE	2.3
1	A	368	PHE	2.3
1	B	557	LYS	2.3
1	B	529	ASN	2.3
1	A	90	TRP	2.3
1	B	575	SER	2.3
1	B	551	ALA	2.3
1	B	458	LEU	2.3
1	B	388	PRO	2.2
1	B	466	GLY	2.2
1	A	563	ILE	2.2
1	B	574	ALA	2.2
1	B	546	LEU	2.2
1	B	495	GLY	2.2
1	B	350	ALA	2.2
1	B	520	THR	2.2
1	B	515	ARG	2.2
1	B	213	LYS	2.1
1	A	683	GLY	2.1
1	B	558	ALA	2.1
1	B	543	LEU	2.1
1	A	25	HIS	2.1
1	A	70	THR	2.1
1	A	95	TYR	2.1
1	B	446	SER	2.1
1	B	585	ALA	2.0
1	B	547	GLY	2.0
1	B	392	ARG	2.0
1	B	449	LEU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.


### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	GLC	A	1742	12/12	0.76	0.30	29,44,48,57	0
4	ACT	A	1743	4/4	0.87	0.15	23,23,24,25	0
3	HEM	B	1741	43/43	0.95	0.14	19,28,34,35	0
3	HEM	A	1741	43/43	0.95	0.15	19,21,29,36	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





## 6.5 Other polymers [i](#)

There are no such residues in this entry.