



# Full wwPDB X-ray Structure Validation Report ⓘ

Sep 5, 2023 – 09:24 PM EDT

PDB ID : 4EN0  
Title : Crystal structure of light  
Authors : Zhan, C.; Liu, W.; Patskovsky, Y.; Ramagopal, U.A.; Bonanno, J.B.; Nathenson, S.G.; Almo, S.C.; New York Structural Genomics Research Consortium (NYSGRG); Atoms-to-Animals: The Immune Function Network (IFN)  
Deposited on : 2012-04-12  
Resolution : 2.59 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.35  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.59 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3163 (2.60-2.60)
Clashscore	141614	3518 (2.60-2.60)
Ramachandran outliers	138981	3455 (2.60-2.60)
Sidechain outliers	138945	3455 (2.60-2.60)
RSRZ outliers	127900	3104 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	167	 10% 66% 19% 13%
1	B	167	 8% 67% 18% 14%
1	C	167	 13% 63% 20% 14%

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 3337 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tumor necrosis factor ligand superfamily member 14.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	145	1103	704	192	204	3	0	0	0
1	B	143	1103	703	193	204	3	0	1	0
1	C	144	1094	697	191	203	3	0	0	0

There are 30 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	74	ARG	-	expression tag	UNP O43557
A	75	SER	-	expression tag	UNP O43557
A	76	HIS	-	expression tag	UNP O43557
A	77	HIS	-	expression tag	UNP O43557
A	78	HIS	-	expression tag	UNP O43557
A	79	HIS	-	expression tag	UNP O43557
A	80	HIS	-	expression tag	UNP O43557
A	81	HIS	-	expression tag	UNP O43557
A	82	GLY	-	expression tag	UNP O43557
A	214	GLU	LYS	SEE REMARK 999	UNP O43557
B	74	ARG	-	expression tag	UNP O43557
B	75	SER	-	expression tag	UNP O43557
B	76	HIS	-	expression tag	UNP O43557
B	77	HIS	-	expression tag	UNP O43557
B	78	HIS	-	expression tag	UNP O43557
B	79	HIS	-	expression tag	UNP O43557
B	80	HIS	-	expression tag	UNP O43557
B	81	HIS	-	expression tag	UNP O43557
B	82	GLY	-	expression tag	UNP O43557
B	214	GLU	LYS	SEE REMARK 999	UNP O43557
C	74	ARG	-	expression tag	UNP O43557
C	75	SER	-	expression tag	UNP O43557
C	76	HIS	-	expression tag	UNP O43557

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	77	HIS	-	expression tag	UNP O43557
C	78	HIS	-	expression tag	UNP O43557
C	79	HIS	-	expression tag	UNP O43557
C	80	HIS	-	expression tag	UNP O43557
C	81	HIS	-	expression tag	UNP O43557
C	82	GLY	-	expression tag	UNP O43557
C	214	GLU	LYS	SEE REMARK 999	UNP O43557

- Molecule 2 is PHOSPHATE ION (three-letter code: PO4) (formula: O<sub>4</sub>P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	P	0	0
			5	4	1		

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula:  $C_8H_{15}NO_6$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			14	8	1	5		


- Molecule 5 is water.

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>	<b>ZeroOcc</b>	<b>AltConf</b>
5	A	6	Total O 6 6	0	0
5	B	3	Total O 3 3	0	0
5	C	3	Total O 3 3	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Tumor necrosis factor ligand superfamily member 14



- Molecule 1: Tumor necrosis factor ligand superfamily member 14



- Molecule 1: Tumor necrosis factor ligand superfamily member 14



## 4 Data and refinement statistics

Property	Value	Source
Space group	I 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	94.70Å 99.98Å 124.38Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.12 – 2.59 46.12 – 2.59	Depositor EDS
% Data completeness (in resolution range)	99.5 (46.12-2.59) 99.3 (46.12-2.59)	Depositor EDS
$R_{merge}$	0.14	Depositor
$R_{sym}$	0.09	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.38 (at 2.58Å)	Xtrriage
Refinement program	PHENIX (phenix.refine: 1.7.3_928)	Depositor
R, $R_{free}$	0.234 , 0.286 0.233 , 0.287	Depositor DCC
$R_{free}$ test set	956 reflections (5.12%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	51.8	Xtrriage
Anisotropy	0.585	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 53.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	3337	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	74.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.77% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, NAG, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.24	0/1129	0.45	0/1533
1	B	0.21	0/1131	0.41	0/1533
1	C	0.32	0/1119	0.47	0/1519
All	All	0.26	0/3379	0.45	0/4585

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1103	0	1074	20	0
1	B	1103	0	1080	19	0
1	C	1094	0	1071	24	0
2	A	5	0	0	1	0
3	A	6	0	8	0	0
4	A	14	0	13	0	0
5	A	6	0	0	0	0
5	B	3	0	0	0	0
5	C	3	0	0	0	0
All	All	3337	0	3246	58	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (58) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:101:ALA:HB3	1:A:230:GLY:HA2	1.64	0.78
1:C:228:ARG:H	1:C:228:ARG:HD2	1.54	0.72
1:A:102:ASN:HB3	1:A:113:LEU:HB2	1.78	0.65
1:C:99:THR:HG22	1:C:230:GLY:HA3	1.79	0.65
1:B:127:SER:HB2	1:B:134:VAL:HB	1.81	0.63
1:B:168:LYS:HB2	1:B:179:LEU:HD11	1.81	0.60
1:A:168:LYS:HB2	1:A:179:LEU:HD11	1.84	0.59
1:C:149:LEU:HD22	1:C:162:ILE:HD13	1.85	0.59
1:C:101:ALA:HB3	1:C:227:LEU:HB3	1.86	0.58
1:C:154:CYS:N	1:C:155:PRO:HD3	2.20	0.56
1:B:98:LEU:HD22	1:B:116:THR:HG22	1.87	0.55
1:A:132:ALA:HB1	1:A:214:GLU:HB3	1.89	0.54
1:A:163:THR:HB	1:A:220:LEU:HB2	1.89	0.54
1:A:134:VAL:HG22	1:A:214:GLU:HG2	1.88	0.53
1:A:105:LEU:HB3	1:A:110:GLY:HA3	1.91	0.53
1:B:152:VAL:HG12	1:B:196:VAL:HG22	1.89	0.53
1:A:228:ARG:NH1	1:C:183:GLN:OE1	2.42	0.52
1:A:151:GLY:HA3	1:A:197:TRP:CE2	2.44	0.52
1:C:168:LYS:HB2	1:C:179:LEU:HD11	1.91	0.51
1:A:98:LEU:HD22	1:A:116:THR:HG22	1.94	0.50
1:C:127:SER:HB2	1:C:134:VAL:HB	1.94	0.50
1:C:158:LEU:HD23	1:C:187:CYS:O	2.12	0.49
1:B:99:THR:HB	1:B:230:GLY:HA3	1.94	0.49
1:B:132:ALA:HB1	1:B:214:GLU:HB3	1.94	0.48
2:A:301:PO4:P	1:B:124:ARG:HH12	2.36	0.48
1:A:175:GLU:H	1:A:175:GLU:HG2	1.55	0.47
1:A:149:LEU:HD22	1:A:162:ILE:HG21	1.97	0.47
1:C:134:VAL:HG22	1:C:214:GLU:HG2	1.95	0.47
1:A:127:SER:HB2	1:A:134:VAL:HB	1.97	0.47
1:C:229:ASP:HB3	1:C:232:ARG:HD3	1.96	0.46
1:A:167:TYR:CE2	1:A:218:ARG:HB2	2.51	0.46
1:B:169[A]:ARG:HB3	1:B:176:GLU:HG2	1.98	0.46
1:C:167:TYR:CE1	1:C:178:GLU:HB3	2.51	0.46
1:A:197:TRP:HB3	1:C:158:LEU:HB3	1.99	0.45
1:A:146:LYS:HD3	1:A:200:SER:OG	2.16	0.45
1:B:162:ILE:HD11	1:B:224:LEU:HD13	1.99	0.45
1:B:229:ASP:OD2	1:B:229:ASP:N	2.49	0.45

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:98:LEU:HB3	1:B:114:TRP:HB3	1.99	0.45
1:C:99:THR:HG22	1:C:100:GLY:H	1.82	0.45
1:C:154:CYS:N	1:C:155:PRO:CD	2.80	0.44
1:A:232:ARG:HG3	1:C:181:VAL:HB	1.99	0.44
1:B:172:ARG:NH1	1:B:173:TYR:OH	2.51	0.43
1:B:93:ASN:O	1:B:124:ARG:NH2	2.51	0.43
1:B:176:GLU:OE2	1:B:218:ARG:NE	2.51	0.43
1:C:94:PRO:HG2	1:C:239:MET:HB2	1.99	0.43
1:A:228:ARG:H	1:A:228:ARG:HG3	1.49	0.42
1:C:104:SER:O	1:C:104:SER:OG	2.27	0.42
1:C:145:SER:HB3	1:C:166:LEU:HD21	2.02	0.42
1:A:190:ALA:HB2	1:C:158:LEU:HD11	2.01	0.42
1:B:169[B]:ARG:HB2	1:B:176:GLU:HG2	2.01	0.42
1:C:167:TYR:CE2	1:C:218:ARG:HB2	2.55	0.41
1:A:181:VAL:HB	1:B:232:ARG:HD2	2.03	0.41
1:B:229:ASP:HB2	1:B:232:ARG:HB2	2.02	0.41
1:B:167:TYR:CE1	1:B:178:GLU:HG2	2.56	0.41
1:C:157:GLY:HA2	1:C:188:GLY:HA2	2.03	0.41
1:B:125:GLY:O	1:B:136:THR:OG1	2.31	0.40
1:C:178:GLU:CD	1:C:181:VAL:HG21	2.40	0.40
1:C:166:LEU:CD2	1:C:217:VAL:HG22	2.52	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	141/167 (84%)	131 (93%)	10 (7%)	0	100	100
1	B	138/167 (83%)	135 (98%)	3 (2%)	0	100	100
1	C	140/167 (84%)	132 (94%)	8 (6%)	0	100	100
All	All	419/501 (84%)	398 (95%)	21 (5%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	113/137 (82%)	104 (92%)	9 (8%)	12	24
1	B	116/137 (85%)	112 (97%)	4 (3%)	37	63
1	C	114/137 (83%)	101 (89%)	13 (11%)	5	10
All	All	343/411 (84%)	317 (92%)	26 (8%)	13	26

All (26) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	99	THR
1	A	146	LYS
1	A	161	THR
1	A	189	ARG
1	A	221	ASP
1	A	222	GLU
1	A	228	ARG
1	A	231	THR
1	A	232	ARG
1	B	175	GLU
1	B	184	GLN
1	B	228	ARG
1	B	229	ASP
1	C	99	THR
1	C	104	SER
1	C	133	LEU
1	C	135	VAL
1	C	137	LYS
1	C	154	CYS
1	C	161	THR
1	C	169	ARG
1	C	177	LEU
1	C	187	CYS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	C	223	ARG
1	C	227	LEU
1	C	229	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	PO4	A	301	-	4,4,4	0.93	0	6,6,6	0.47	0
3	GOL	A	302	-	5,5,5	0.37	0	5,5,5	0.28	0
4	NAG	A	303	1	14,14,15	0.54	0	17,19,21	1.22	2 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	302	-	-	2/4/4/4	-
4	NAG	A	303	1	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	303	NAG	C1-O5-C5	3.52	116.97	112.19
4	A	303	NAG	C4-C3-C2	-2.01	108.07	111.02

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	302	GOL	O1-C1-C2-C3
3	A	302	GOL	O1-C1-C2-O2

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	301	PO4	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	145/167 (86%)	0.79	17 (11%) <b>4</b> <b>3</b>	36, 69, 108, 135	0
1	B	143/167 (85%)	0.58	14 (9%) <b>7</b> <b>5</b>	38, 66, 112, 146	0
1	C	144/167 (86%)	1.05	22 (15%) <b>2</b> <b>1</b>	48, 77, 122, 149	0
All	All	432/501 (86%)	0.80	53 (12%) <b>4</b> <b>2</b>	36, 71, 117, 149	0

All (53) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	103	SER	6.4
1	A	155	PRO	6.3
1	B	190	ALA	5.8
1	C	194	SER	5.7
1	C	104	SER	5.6
1	A	157	GLY	5.3
1	C	190	ALA	4.7
1	C	227	LEU	4.5
1	A	172	ARG	4.1
1	A	171	PRO	4.0
1	C	99	THR	3.8
1	C	173	TYR	3.8
1	B	191	THR	3.8
1	B	155	PRO	3.8
1	B	104	SER	3.7
1	A	191	THR	3.7
1	C	222	GLU	3.6
1	B	156	LEU	3.5
1	B	172	ARG	3.5
1	C	193	SER	3.4
1	C	118	LEU	3.4
1	C	192	SER	3.4
1	A	228	ARG	3.4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	173	TYR	3.4
1	A	211	ALA	3.3
1	B	92	VAL	3.3
1	A	190	ALA	3.2
1	C	189	ARG	3.1
1	B	125	GLY	3.1
1	A	158	LEU	3.1
1	C	171	PRO	3.1
1	C	181	VAL	3.0
1	A	152	VAL	2.9
1	C	117	GLN	2.9
1	C	223	ARG	2.9
1	B	211	ALA	2.8
1	B	189	ARG	2.7
1	A	93	ASN	2.6
1	C	172	ARG	2.6
1	A	125	GLY	2.4
1	C	155	PRO	2.4
1	A	232	ARG	2.4
1	A	231	THR	2.3
1	A	160	SER	2.3
1	A	129	HIS	2.3
1	B	108	SER	2.2
1	C	235	PHE	2.2
1	C	95	ALA	2.1
1	B	192	SER	2.1
1	C	191	THR	2.1
1	B	138	ALA	2.1
1	C	240	VAL	2.1
1	A	101	ALA	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.



## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	NAG	A	303	14/15	0.86	0.15	68,83,96,96	0
3	GOL	A	302	6/6	0.92	0.49	78,87,91,99	0
2	PO4	A	301	5/5	0.92	0.21	131,131,134,140	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.