



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 23, 2024 – 08:08 AM EDT

PDB ID : 4RHS
Title : Crystal structure of GD2 bound PltB
Authors : Gao, X.; Wang, J.; Galan, J.
Deposited on : 2014-10-02
Resolution : 1.92 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.37.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.37.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION






The reported resolution of this entry is 1.92 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	7937 (1.94-1.90)
Clashscore	141614	8644 (1.94-1.90)
Ramachandran outliers	138981	8530 (1.94-1.90)
Sidechain outliers	138945	8530 (1.94-1.90)
RSRZ outliers	127900	7793 (1.94-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	122	
1	B	122	
1	C	122	
1	D	122	
1	E	122	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain	
2	F	3	 33%	67%
2	G	3	 33%	67%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4962 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Putative pertussis-like toxin subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	114	885	566	145	170	4	0	0	0
1	B	115	890	569	146	171	4	0	0	0
1	C	114	885	566	145	170	4	0	0	0
1	D	114	885	566	145	170	4	0	0	0
1	E	114	885	566	145	170	4	0	0	0

There are 40 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	138	LEU	-	expression tag	UNP Q8Z6A3
A	139	GLU	-	expression tag	UNP Q8Z6A3
A	140	HIS	-	expression tag	UNP Q8Z6A3
A	141	HIS	-	expression tag	UNP Q8Z6A3
A	142	HIS	-	expression tag	UNP Q8Z6A3
A	143	HIS	-	expression tag	UNP Q8Z6A3
A	144	HIS	-	expression tag	UNP Q8Z6A3
A	145	HIS	-	expression tag	UNP Q8Z6A3
B	138	LEU	-	expression tag	UNP Q8Z6A3
B	139	GLU	-	expression tag	UNP Q8Z6A3
B	140	HIS	-	expression tag	UNP Q8Z6A3
B	141	HIS	-	expression tag	UNP Q8Z6A3
B	142	HIS	-	expression tag	UNP Q8Z6A3
B	143	HIS	-	expression tag	UNP Q8Z6A3
B	144	HIS	-	expression tag	UNP Q8Z6A3
B	145	HIS	-	expression tag	UNP Q8Z6A3
C	138	LEU	-	expression tag	UNP Q8Z6A3
C	139	GLU	-	expression tag	UNP Q8Z6A3
C	140	HIS	-	expression tag	UNP Q8Z6A3

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	141	HIS	-	expression tag	UNP Q8Z6A3
C	142	HIS	-	expression tag	UNP Q8Z6A3
C	143	HIS	-	expression tag	UNP Q8Z6A3
C	144	HIS	-	expression tag	UNP Q8Z6A3
C	145	HIS	-	expression tag	UNP Q8Z6A3
D	138	LEU	-	expression tag	UNP Q8Z6A3
D	139	GLU	-	expression tag	UNP Q8Z6A3
D	140	HIS	-	expression tag	UNP Q8Z6A3
D	141	HIS	-	expression tag	UNP Q8Z6A3
D	142	HIS	-	expression tag	UNP Q8Z6A3
D	143	HIS	-	expression tag	UNP Q8Z6A3
D	144	HIS	-	expression tag	UNP Q8Z6A3
D	145	HIS	-	expression tag	UNP Q8Z6A3
E	138	LEU	-	expression tag	UNP Q8Z6A3
E	139	GLU	-	expression tag	UNP Q8Z6A3
E	140	HIS	-	expression tag	UNP Q8Z6A3
E	141	HIS	-	expression tag	UNP Q8Z6A3
E	142	HIS	-	expression tag	UNP Q8Z6A3
E	143	HIS	-	expression tag	UNP Q8Z6A3
E	144	HIS	-	expression tag	UNP Q8Z6A3
E	145	HIS	-	expression tag	UNP Q8Z6A3

- Molecule 2 is an oligosaccharide called N-acetyl-alpha-neuraminic acid-(2-8)-N-acetyl-alpha-neuraminic acid-(2-3)-beta-D-galactopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	F	3	52	28	2	22	0	0	0
2	G	3	52	28	2	22	0	0	0

- Molecule 3 is ACETATE ION (three-letter code: ACT) (formula: C₂H₃O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	1	Total C O 4 2 2	0	0
3	B	1	Total C O 4 2 2	0	0



- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	104	Total O 104 104	0	0
4	B	108	Total O 108 108	0	0
4	C	72	Total O 72 72	0	0
4	D	56	Total O 56 56	0	0
4	E	80	Total O 80 80	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: Putative pertussis-like toxin subunit





- Molecule 1: Putative pertussis-like toxin subunit





- Molecule 1: Putative pertussis-like toxin subunit



- Molecule 1: Putative pertussis-like toxin subunit



- Molecule 1: Putative pertussis-like toxin subunit



- Molecule 2: N-acetyl-alpha-neuraminic acid-(2-8)-N-acetyl-alpha-neuraminic acid-(2-3)-beta-D-galactopyranose

Chain F:  33% 67%

CA11
SIA2
SIA3

- Molecule 2: N-acetyl-alpha-neuraminic acid-(2-8)-N-acetyl-alpha-neuraminic acid-(2-3)-beta-D-galactopyranose

Chain G:  33% 67%

CA11
SIA2
SIA3

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	68.26Å 96.75Å 104.00Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	26.05 – 1.92 26.05 – 1.92	Depositor EDS
% Data completeness (in resolution range)	95.6 (26.05-1.92) 95.6 (26.05-1.92)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.06 (at 1.92Å)	Xtrriage
Refinement program	PHENIX (phenix.refine: dev_1701)	Depositor
R, R_{free}	0.175 , 0.205 0.179 , 0.206	Depositor DCC
R_{free} test set	2571 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	25.9	Xtrriage
Anisotropy	0.397	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 54.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4962	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.71% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SIA, GAL, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.40	0/909	0.54	0/1240
1	B	0.42	0/914	0.57	0/1247
1	C	0.42	0/909	0.56	0/1240
1	D	0.36	0/909	0.53	0/1240
1	E	0.38	0/909	0.55	0/1240
All	All	0.40	0/4550	0.55	0/6207

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	885	0	846	4	0
1	B	890	0	848	8	0
1	C	885	0	846	6	0
1	D	885	0	846	3	0
1	E	885	0	846	3	0
2	F	52	0	44	0	0
2	G	52	0	44	0	0
3	B	8	0	6	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	104	0	0	2	1
4	B	108	0	0	3	1
4	C	72	0	0	1	0
4	D	56	0	0	0	0
4	E	80	0	0	2	0
All	All	4962	0	4326	22	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (22) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:50:SER:OG	1:D:73:SER:O	2.07	0.72
1:C:100:ARG:HH12	1:C:137:LYS:HB3	1.61	0.65
1:C:100:ARG:NH1	1:C:137:LYS:HB3	2.24	0.52
1:B:84:GLU:OE2	4:B:369:HOH:O	2.20	0.49
1:A:128:CYS:O	4:A:218:HOH:O	2.19	0.48
1:A:137:LYS:NZ	4:A:232:HOH:O	2.37	0.47
1:B:36:ASP:HA	1:B:98:SER:OG	2.16	0.46
1:C:53:PHE:CE1	1:C:71:ALA:HB3	2.51	0.46
1:D:128:CYS:HA	1:D:133:CYS:HA	1.96	0.46
1:A:81:SER:OG	1:C:46:GLN:HB2	2.17	0.45
1:B:128:CYS:HA	1:B:133:CYS:HA	1.99	0.44
1:C:81:SER:OG	1:D:46:GLN:HB2	2.18	0.44
1:E:100:ARG:HH12	1:E:137:LYS:HB3	1.84	0.43
1:A:100:ARG:CZ	1:A:133:CYS:HB2	2.50	0.42
1:C:137:LYS:NZ	4:C:354:HOH:O	2.40	0.42
1:E:96:GLY:HA2	4:E:302:HOH:O	2.19	0.42
1:B:24:GLU:N	1:B:24:GLU:OE1	2.54	0.41
1:B:83:LYS:NZ	4:B:381:HOH:O	2.54	0.41
1:E:90:ARG:HD2	4:E:365:HOH:O	2.21	0.41
1:B:47:ILE:HD11	3:B:201:ACT:H3	2.03	0.40
1:B:62:GLY:HA2	4:B:342:HOH:O	2.20	0.40
1:B:110:TYR:HA	1:B:111:PRO:HD3	1.87	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:239:HOH:O	4:B:330:HOH:O[3_544]	2.07	0.13

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	112/122 (92%)	109 (97%)	3 (3%)	0	100	100
1	B	113/122 (93%)	110 (97%)	3 (3%)	0	100	100
1	C	112/122 (92%)	104 (93%)	8 (7%)	0	100	100
1	D	112/122 (92%)	108 (96%)	4 (4%)	0	100	100
1	E	112/122 (92%)	110 (98%)	2 (2%)	0	100	100
All	All	561/610 (92%)	541 (96%)	20 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	98/106 (92%)	98 (100%)	0	100	100
1	B	98/106 (92%)	97 (99%)	1 (1%)	76	75
1	C	98/106 (92%)	96 (98%)	2 (2%)	55	49
1	D	98/106 (92%)	97 (99%)	1 (1%)	76	75
1	E	98/106 (92%)	98 (100%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
All	All	490/530 (92%)	486 (99%)	4 (1%)	81	81

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	110	TYR
1	C	98	SER
1	C	110	TYR
1	D	76	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

6 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GAL	F	1	2	12,12,12	0.80	0	17,17,17	0.94	0
2	SIA	F	2	2	20,20,21	2.22	2 (10%)	24,28,31	1.78	4 (16%)
2	SIA	F	3	2	20,20,21	2.10	2 (10%)	24,28,31	1.49	4 (16%)
2	GAL	G	1	2	12,12,12	0.73	0	17,17,17	0.79	0
2	SIA	G	2	2	20,20,21	2.20	2 (10%)	24,28,31	1.62	4 (16%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	SIA	G	3	2	20,20,21	1.99	2 (10%)	24,28,31	1.58	4 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GAL	F	1	2	-	2/2/22/22	0/1/1/1
2	SIA	F	2	2	-	3/18/34/38	0/1/1/1
2	SIA	F	3	2	-	4/18/34/38	0/1/1/1
2	GAL	G	1	2	-	1/2/22/22	0/1/1/1
2	SIA	G	2	2	-	2/18/34/38	0/1/1/1
2	SIA	G	3	2	-	5/18/34/38	0/1/1/1

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	F	2	SIA	C2-C1	8.58	1.60	1.52
2	G	2	SIA	C2-C1	8.39	1.59	1.52
2	F	3	SIA	C2-C1	8.15	1.59	1.52
2	G	3	SIA	C2-C1	7.77	1.59	1.52
2	F	2	SIA	O6-C2	2.85	1.47	1.43
2	F	3	SIA	O6-C2	2.76	1.47	1.43
2	G	2	SIA	O6-C2	2.76	1.47	1.43
2	G	3	SIA	O6-C2	2.26	1.46	1.43

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	F	2	SIA	O6-C2-C3	-4.52	104.24	110.46
2	F	3	SIA	O1A-C1-C2	-4.17	112.72	122.57
2	G	2	SIA	O6-C2-C3	-3.91	105.08	110.46
2	F	2	SIA	O1A-C1-C2	-3.80	113.60	122.57
2	F	2	SIA	C6-O6-C2	3.78	119.43	111.34
2	G	3	SIA	O1A-C1-C2	-3.64	113.98	122.57
2	G	2	SIA	C6-O6-C2	3.45	118.73	111.34
2	G	2	SIA	O1A-C1-C2	-3.30	114.77	122.57
2	G	3	SIA	C6-O6-C2	3.23	118.24	111.34
2	F	3	SIA	C6-O6-C2	2.80	117.33	111.34
2	G	3	SIA	O1B-C1-O1A	2.60	130.00	124.09

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	F	3	SIA	O1B-C1-O1A	2.59	129.98	124.09
2	G	2	SIA	O1B-C1-O1A	2.18	129.04	124.09
2	G	3	SIA	O6-C2-C1	2.12	111.86	107.70
2	F	3	SIA	C8-C7-C6	-2.12	109.01	113.03
2	F	2	SIA	O4-C4-C3	2.09	115.12	109.94

There are no chirality outliers.


All (17) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	F	2	SIA	C7-C8-C9-O9
2	G	3	SIA	C6-C7-C8-C9
2	G	3	SIA	C6-C7-C8-O8
2	G	3	SIA	O7-C7-C8-C9
2	G	3	SIA	O7-C7-C8-O8
2	F	2	SIA	O8-C8-C9-O9
2	F	3	SIA	O7-C7-C8-O8
2	F	1	GAL	O5-C5-C6-O6
2	F	3	SIA	C6-C7-C8-O8
2	F	3	SIA	O7-C7-C8-C9
2	F	3	SIA	C6-C7-C8-C9
2	G	2	SIA	O8-C8-C9-O9
2	F	1	GAL	C4-C5-C6-O6
2	G	3	SIA	C7-C8-C9-O9
2	G	2	SIA	C7-C8-C9-O9
2	F	2	SIA	O1A-C1-C2-O6
2	G	1	GAL	C4-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	ACT	B	202	-	3,3,3	0.72	0	3,3,3	1.32	0
3	ACT	B	201	-	3,3,3	0.79	0	3,3,3	1.38	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	201	ACT	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	114/122 (93%)	0.02	3 (2%) 56 59	17, 24, 40, 65	0
1	B	115/122 (94%)	-0.08	2 (1%) 70 72	18, 24, 46, 54	0
1	C	114/122 (93%)	0.33	14 (12%) 4 4	18, 27, 61, 97	0
1	D	114/122 (93%)	0.53	14 (12%) 4 4	23, 34, 59, 74	0
1	E	114/122 (93%)	0.39	10 (8%) 10 11	19, 28, 57, 71	0
All	All	571/610 (93%)	0.24	43 (7%) 14 16	17, 27, 53, 97	0

All (43) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	130	ALA	5.4
1	C	131	THR	4.8
1	A	131	THR	4.7
1	D	29	ASN	4.6
1	D	65	THR	4.2
1	E	111	PRO	3.9
1	E	109	THR	3.5
1	C	63	SER	3.3
1	E	130	ALA	3.3
1	E	110	TYR	3.1
1	C	62	GLY	3.1
1	C	30	THR	3.1
1	A	130	ALA	3.1
1	C	29	ASN	3.0
1	C	61	ASN	3.0
1	E	131	THR	2.8
1	D	131	THR	2.8
1	C	65	THR	2.8
1	E	115	ASN	2.7
1	B	65	THR	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	132	GLN	2.6
1	C	122	LEU	2.6
1	D	130	ALA	2.5
1	D	64	GLY	2.5
1	D	30	THR	2.5
1	E	112	LEU	2.5
1	A	112	LEU	2.4
1	C	125	LEU	2.3
1	D	122	LEU	2.3
1	D	110	TYR	2.3
1	E	55	ILE	2.2
1	D	28	ASP	2.2
1	D	111	PRO	2.2
1	D	112	LEU	2.2
1	C	137	LYS	2.2
1	D	137	LYS	2.2
1	C	64	GLY	2.2
1	C	91	TYR	2.2
1	D	106	ASN	2.1
1	D	109	THR	2.1
1	E	44	VAL	2.1
1	B	29	ASN	2.1
1	E	64	GLY	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	GAL	F	1	12/12	0.74	0.35	75,78,83,89	0
2	GAL	G	1	12/12	0.81	0.28	57,69,73,78	0
2	SIA	F	2	20/21	0.84	0.27	46,57,71,72	0
2	SIA	F	3	20/21	0.91	0.19	25,48,57,58	0
2	SIA	G	2	20/21	0.92	0.19	28,43,62,62	0


Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	SIA	G	3	20/21	0.95	0.18	27,36,53,55	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	ACT	B	201	4/4	0.93	0.16	33,44,45,51	0
3	ACT	B	202	4/4	0.95	0.10	31,36,40,41	0

6.5 Other polymers [i](#)

There are no such residues in this entry.