

# Full wwPDB X-ray Structure Validation Report (i)

Dec 16, 2023 – 11:13 am GMT

PDB ID : 4TTI

Title : Crystal structure of double mutant E. Coli purine nucleoside phosphorylase

with 4 FMC molecules

Authors : Stefanic, Z.; Bzowska, A.

Deposited on : 2014-06-21

Resolution : 1.89 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul : 1.8.4, CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.36

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

 $Refmac \quad : \quad 5.8.0158$ 

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.36

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 1.89 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	Similar resolution $(\# \text{Entries, resolution range}(\text{\AA}))$
$R_{free}$	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	A	237	95%	•
1	В	237	90%	9% •
1	С	237	85%	8% • 5%
1	D	237	96%	
1	Е	237	89%	11%



 $Continued\ from\ previous\ page...$ 

Mol	Chain	Length	Quality of chain			
1	F	237	95%	5%		



# 2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 12099 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Purine nucleoside phosphorylase DeoD-type.


Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
1	A	237	Total	С	N	О	S	0	3	0
1	A	231	1809	1145	311	337	16	U	3	
1	В	237	Total	С	N	О	S	0	3	0
1	Ъ	231	1805	1144	308	337	16	U	J	
1	С	224	Total	С	N	О	S	0	3	0
1		224	1701	1082	286	317	16	U		
1	D	237	Total	С	N	О	S	0	1	0
1	D	231	1793	1134	307	337	15	U	1	
1	Е	237	Total	С	N	Ο	S	0	1	0
1	ш	201	1793	1135	305	337	16	0	1	
1	F	237	Total	С	N	О	S	0	4	0
1	I'	231	1820	1151	317	337	15	U	4	U

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	204	ALA	ASP	engineered mutation	UNP P0ABP8
A	217	ALA	ARG	engineered mutation	UNP P0ABP8
В	204	ALA	ASP	engineered mutation	UNP P0ABP8
В	217	ALA	ARG	engineered mutation	UNP P0ABP8
С	204	ALA	ASP	engineered mutation	UNP P0ABP8
С	217	ALA	ARG	engineered mutation	UNP P0ABP8
D	204	ALA	ASP	engineered mutation	UNP P0ABP8
D	217	ALA	ARG	engineered mutation	UNP P0ABP8
Е	204	ALA	ASP	engineered mutation	UNP P0ABP8
Е	217	ALA	ARG	engineered mutation	UNP P0ABP8
F	204	ALA	ASP	engineered mutation	UNP P0ABP8
F	217	ALA	ARG	engineered mutation	UNP P0ABP8


• Molecule 2 is (1S)-1-(7-amino-1H-pyrazolo[4,3-d]pyrimidin-3-yl)-1,4-anhydro-D-ribitol (three-letter code: FMC) (formula:  $C_{10}H_{13}N_5O_4$ ).





Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	Λ	1	Total	С	N	О	0	0
2	A	1	19	10	5	4	0	
2	C	1	Total	С	N	О	0	0
2		1	19	10	5	4	0	0
2	D	1	Total	С	N	О	0	0
2	ט	1	19	10	5	4	0	0
2	F	1	Total	С	N	О	0	0
	I'	1	19	10	5	4	0	0


 $\bullet$  Molecule 3 is PHOSPHATE ION (three-letter code: PO4) (formula:  $\mathrm{O_4P}).$ 





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total O P 5 4 1	0	0
3	В	1	Total O P 5 4 1	0	0
3	С	1	Total O P 5 4 1	0	0
3	D	1	Total O P 5 4 1	0	0
3	E	1	Total O P 5 4 1	0	0
3	F	1	Total O P 5 4 1	0	0

 $\bullet$  Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	С	1	Total O S 5 4 1	0	0
4	С	1	Total O S 5 4 1	0	0
4	D	1	Total O S 5 4 1	0	0
4	E	1	Total O S 5 4 1	0	0
4	F	1	Total O S 5 4 1	0	0

• Molecule 5 is water.




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	237	Total O 237 237	0	0
5	В	183	Total O 183 183	9	0
5	С	206	Total O 206 206	0	0
5	D	192	Total O 192 192	0	0
5	Е	193	Total O 193 193	6	0
5	F	236	Total O 236 236	0	0



# 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Purine nucleoside phosphorylase DeoD-type



• Molecule 1: Purine nucleoside phosphorylase DeoD-type



Chain F: 95% 5%



# 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	61.93Å 124.17Å 189.44Å	Donogitor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.70 - 1.89	Depositor
Resolution (A)	47.83 - 1.89	EDS
% Data completeness	98.8 (47.70-1.89)	Depositor
(in resolution range)	98.8 (47.83-1.89)	EDS
$R_{merge}$	0.17	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) > 1$	1.44 (at 1.90Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.2_1309)	Depositor
P. P.	0.167 , 0.210	Depositor
$R, R_{free}$	0.166 , 0.210	DCC
$R_{free}$ test set	5807  reflections  (5.00%)	wwPDB-VP
Wilson B-factor $(\mathring{A}^2)$	17.7	Xtriage
Anisotropy	0.400	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$ , $B_{sol}(Å^2)$	0.37, 51.5	EDS
L-test for twinning <sup>2</sup>	$  <  L  > = 0.48, < L^2 > = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	12099	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 3.17% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of <|L|>,  $<L^2>$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

# 5 Model quality (i)

## 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, FMC, PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bo	nd lengths	Bond angles	
Moi   Chain		RMSZ	RMSZ $ $ # $ Z  > 5$		# Z >5
1	A	0.37	0/1847	0.52	0/2488
1	В	0.37	0/1843	0.53	0/2484
1	С	0.36	0/1737	0.55	0/2339
1	D	0.36	0/1825	0.53	0/2461
1	Е	0.36	0/1825	0.53	0/2460
1	F	0.41	1/1861 (0.1%)	0.53	0/2506
All	All	0.37	1/10938 (0.0%)	0.53	0/14738

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a maintain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	С	0	1

#### All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\text{\AA})$	Ideal(Å)
1	F	216	GLU	CD-OE2	7.17	1.33	1.25

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	С	219	THR	Peptide



### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1809	0	1830	4	0
1	В	1805	0	1826	16	0
1	С	1701	0	1721	15	0
1	D	1793	0	1799	5	0
1	Ε	1793	0	1804	14	0
1	F	1820	0	1847	7	0
2	A	19	0	13	1	0
2	С	19	0	13	2	0
2	D	19	0	13	1	0
2	F	19	0	13	2	0
3	A	5	0	0	0	0
3	В	5	0	0	0	0
3	С	5	0	0	1	0
3	D	5	0	0	0	0
3	Е	5	0	0	0	0
3	F	5	0	0	0	0
4	С	10	0	0	1	0
4	D	5	0	0	0	0
4	Е	5	0	0	1	0
4	F	5	0	0	0	0
5	A	237	0	0	1	0
5	В	183	0	0	2	0
5	С	206	0	0	2	0
5	D	192	0	0	0	0
5	Ε	193	0	0	2	0
5	F	236	0	0	5	0
All	All	12099	0	10879	61	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (61) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:219:THR:HB	1:C:221:PHE:H	1.39	0.87



 $Continued\ from\ previous\ page...$ 

Continuea from previo		Interatomic	Clash
Atom-1	Atom-2	distance (Å)	$\text{overlap } (\mathring{\mathrm{A}})$
1:C:219:THR:HB	1:C:221:PHE:N	2.05	0.70
1:C:222:ASN:ND2	4:C:303:SO4:O2	2.23	0.68
1:C:219:THR:HA	1:C:220:THR:HB	1.79	0.64
1:C:237:LYS:O	5:C:557:HOH:O	2.15	0.63
1:E:95:LEU:HD23	1:E:171:GLU:HG3	1.81	0.62
1:E:35:ASP:O	5:E:549:HOH:O	2.17	0.59
1:F:30:GLU:OE1	5:F:401:HOH:O	2.17	0.59
2:D:302:FMC:N3	2:D:302:FMC:H2'	2.18	0.58
1:B:101:ARG:HB3	1:B:220:THR:HG21	1.85	0.58
1:B:117:ARG:NH2	1:D:122:ASP:O	2.35	0.57
1:B:30:GLU:OE1	5:B:401:HOH:O	2.17	0.56
1:F:37[B]:ARG:NH1	5:F:612:HOH:O	2.38	0.56
1:F:37[B]:ARG:NH1	5:F:616:HOH:O	2.40	0.55
2:A:300:FMC:N3	2:A:300:FMC:H2'	2.22	0.54
1:E:220:THR:HG21	4:E:302:SO4:O1	2.08	0.53
1:A:57:ILE:HD11	1:A:232:VAL:HG11	1.91	0.53
1:E:15:VAL:HG22	1:E:84:LYS:HB2	1.90	0.53
1:F:205:HIS:CE1	1:F:207:ARG:HB2	2.43	0.53
1:B:100:LEU:HA	1:B:202:VAL:HG22	1.90	0.53
1:C:97:HIS:O	1:C:149:ARG:NH1	2.41	0.53
2:C:301:FMC:H2'	2:C:301:FMC:N3	2.23	0.53
1:C:24:ARG:HD3	1:C:221:PHE:CE1	2.45	0.52
1:B:101:ARG:HE	1:B:220:THR:CG2	2.23	0.51
1:E:220:THR:HG23	1:E:223:ASP:H	1.75	0.51
1:B:114:LYS:HD3	1:E:114:LYS:HG3	1.93	0.51
1:E:13:ALA:HB2	1:E:56:LYS:HG2	1.92	0.50
1:A:236:ASP:O	1:A:237:LYS:HB2	2.11	0.49
1:A:13:ALA:HB2	1:A:56:LYS:HG2	1.95	0.49
2:F:301:FMC:H2'	2:F:301:FMC:N3	2.28	0.49
1:E:226:LYS:NZ	5:E:586:HOH:O	2.45	0.48
1:B:212:THR:OG1	1:B:216:GLU:HG3	2.13	0.48
1:F:149[A]:ARG:NE	5:F:566:HOH:O	2.47	0.48
1:C:62:HIS:CE1	1:C:181:GLU:HG2	2.48	0.48
1:E:220:THR:HG22	1:E:223:ASP:CG	2.34	0.47
5:B:405:HOH:O	1:D:172:LYS:HG3	2.13	0.47
1:E:220:THR:HG23	1:E:222:ASN:N	2.30	0.46
1:D:62:HIS:CE1	1:D:181:GLU:HG2	2.50	0.46
1:B:73:THR:HG22	1:B:85:ILE:HD13	1.97	0.46
1:C:29:ALA:HA	1:C:33:LEU:HD22	1.98	0.46
1:B:43:ARG:HA	1:E:21:ASP:OD1	2.16	0.45
1:E:220:THR:CG2	1:E:223:ASP:H	2.29	0.45



Continued from previous page...

Atom-1	Atom-2	$\begin{array}{c} \text{Interatomic} \\ \text{distance } (\text{\AA}) \end{array}$	Clash overlap (Å)
2:C:301:FMC:H1'	3:C:302:PO4:O2	2.17	0.45
2:F:301:FMC:N7	5:F:599:HOH:O	2.33	0.44
1:C:73:THR:HG22	1:C:85:ILE:HD13	1.99	0.44
1:D:172:LYS:HG2	1:D:173:TYR:CD1	2.53	0.44
1:E:76:LEU:HB3	1:E:82:VAL:HG21	1.99	0.43
1:B:107[A]:MET:HE2	1:B:107[A]:MET:HB2	1.84	0.43
1:C:173:TYR:CZ	1:E:119:ARG:HB3	2.52	0.43
1:B:212:THR:HG23	1:B:217:ALA:HB2	2.00	0.43
1:C:83:LYS:NZ	5:C:548:HOH:O	2.47	0.43
1:B:74:LYS:HD3	1:B:74:LYS:C	2.39	0.43
1:C:43:ARG:HA	1:F:21:ASP:OD1	2.20	0.42
1:A:41:ASN:OD1	5:A:583:HOH:O	2.22	0.42
1:C:90:SER:HB2	1:C:203:SER:HB3	2.02	0.42
1:B:101:ARG:HE	1:B:220:THR:HG21	1.84	0.42
1:C:142:LYS:HE3	1:C:142:LYS:HB2	1.85	0.41
1:B:105:ILE:HG23	1:B:197:LEU:HD11	2.02	0.41
1:B:95:LEU:HD23	1:B:171:GLU:HG3	2.03	0.41
1:F:62:HIS:CE1	1:F:181:GLU:HG2	2.55	0.41
1:B:128:ILE:HD12	1:D:107:MET:HE3	2.02	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	A	$238/237 \ (100\%)$	232 (98%)	6 (2%)	0	100	100
1	В	$238/237 \ (100\%)$	232 (98%)	6 (2%)	0	100	100
1	С	223/237~(94%)	216 (97%)	6 (3%)	1 (0%)	34	24
1	D	$236/237 \ (100\%)$	225 (95%)	9 (4%)	2 (1%)	19	9
1	E	$236/237 \ (100\%)$	225 (95%)	9 (4%)	2 (1%)	19	9



Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	F	$239/237 \ (101\%)$	232 (97%)	7 (3%)	0	100	100
All	All	1410/1422 (99%)	1362 (97%)	43 (3%)	5 (0%)	34	24

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	209	HIS
1	D	215	ALA
1	Е	212	THR
1	Е	215	ALA
1	С	220	THR

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Perce	ntiles
1	A	$189/186\ (102\%)$	183 (97%)	6 (3%)	39	30
1	В	189/186~(102%)	183 (97%)	6 (3%)	39	30
1	С	179/186 (96%)	175 (98%)	4 (2%)	52	47
1	D	186/186 (100%)	183 (98%)	3 (2%)	62	60
1	E	$187/186\ (100\%)$	180 (96%)	7 (4%)	34	25
1	F	$190/186\ (102\%)$	188 (99%)	2 (1%)	73	73
All	All	1120/1116 (100%)	1092 (98%)	28 (2%)	49	41

All (28) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	37	ARG
1	A	107[A]	MET
1	A	107[B]	MET
1	A	179	GLU
1	A	210	GLU
1	A	237	LYS



Continued from previous page...

Mol	Chain	Res	Type
1	В	179	GLU
1	В	202	VAL
1	В	207	ARG
1	В	208	THR
1	В	211	GLN
1	В	212	THR
1	C C C	33	LEU
1	С	101	ARG
1	С	149	ARG
1		179	GLU
1	D	107	MET
1	D	179	GLU
1	D	209	HIS
1	Е	23	LEU
1	Е	34	GLU
1	Е	37	ARG
1	Е	100	LEU
1	Е	120	PHE
1	Е	179	GLU
1	Е	210	GLU
1	F	55	ARG
1	F	179	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	211	GLN

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates (i)

There are no monosaccharides in this entry.



## 5.6 Ligand geometry (i)

15 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

N / - 1	Т	Clasica	Das	Link	Bo	ond leng	ths	В	ond ang	eles
Mol	Type	Chain	Res	Lilik	Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	PO4	С	302	-	4,4,4	0.95	0	6,6,6	0.51	0
4	SO4	С	304	-	4,4,4	0.15	0	6,6,6	0.09	0
2	FMC	A	300	-	20,21,21	1.35	2 (10%)	19,31,31	1.99	3 (15%)
4	SO4	F	303	-	4,4,4	0.15	0	6,6,6	0.18	0
3	PO4	A	301	-	4,4,4	1.22	0	6,6,6	0.80	0
4	SO4	С	303	-	4,4,4	0.13	0	6,6,6	0.14	0
3	PO4	В	301	-	4,4,4	0.84	0	6,6,6	0.60	0
2	FMC	F	301	-	20,21,21	1.35	3 (15%)	19,31,31	1.93	4 (21%)
3	PO4	D	301	-	4,4,4	0.91	0	6,6,6	0.36	0
3	PO4	Е	301	-	4,4,4	0.89	0	6,6,6	0.57	0
4	SO4	D	303	-	4,4,4	0.15	0	6,6,6	0.19	0
2	FMC	D	302	-	20,21,21	1.37	2 (10%)	19,31,31	2.07	3 (15%)
2	FMC	С	301	-	20,21,21	1.27	1 (5%)	19,31,31	1.94	4 (21%)
4	SO4	Е	302	-	4,4,4	0.15	0	6,6,6	0.22	0
3	PO4	F	302	_	4,4,4	1.14	0	6,6,6	0.38	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FMC	D	302	-	-	1/2/22/22	0/3/3/3
2	FMC	С	301	-	-	0/2/22/22	0/3/3/3
2	FMC	A	300	-	-	0/2/22/22	0/3/3/3
2	FMC	F	301	-	-	2/2/22/22	0/3/3/3

All (8) bond length outliers are listed below:



Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	$Ideal(\AA)$
2	D	302	FMC	C9-C4	4.17	1.50	1.43
2	A	300	FMC	C9-C4	3.93	1.49	1.43
2	F	301	FMC	C9-C4	3.80	1.49	1.43
2	С	301	FMC	C9-C4	3.67	1.49	1.43
2	A	300	FMC	N7-N8	2.26	1.42	1.37
2	F	301	FMC	C2-N3	2.21	1.35	1.32
2	D	302	FMC	N7-N8	2.18	1.41	1.37
2	F	301	FMC	N7-N8	2.02	1.41	1.37

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	$Observed(^o)$	$Ideal(^{o})$
2	D	302	FMC	C2-N3-C4	6.44	119.82	114.81
2	A	300	FMC	C2-N3-C4	5.95	119.44	114.81
2	С	301	FMC	C2-N3-C4	5.71	119.25	114.81
2	F	301	FMC	C2-N3-C4	5.46	119.06	114.81
2	D	302	FMC	C5-C4-N3	-4.14	120.10	124.92
2	A	300	FMC	C5-C4-N3	-3.96	120.32	124.92
2	F	301	FMC	C5-C4-N3	-3.90	120.39	124.92
2	С	301	FMC	C5-C4-N3	-3.55	120.80	124.92
2	С	301	FMC	N3-C2-N1	-3.17	123.72	128.68
2	D	302	FMC	N3-C2-N1	-3.04	123.93	128.68
2	F	301	FMC	N3-C2-N1	-2.98	124.01	128.68
2	A	300	FMC	N3-C2-N1	-2.95	124.07	128.68
2	С	301	FMC	C2-N1-C6	2.17	122.47	118.75
2	F	301	FMC	C2-N1-C6	2.02	122.21	118.75

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	F	301	FMC	O4'-C4'-C5'-O5'
2	F	301	FMC	C3'-C4'-C5'-O5'
2	D	302	FMC	O4'-C4'-C5'-O5'

There are no ring outliers.

7 monomers are involved in 8 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	С	302	PO4	1	0
2	A	300	FMC	1	0
4	С	303	SO4	1	0




Continued from previous page...


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	F	301	FMC	2	0
2	D	302	FMC	1	0
2	С	301	FMC	2	0
4	Е	302	SO4	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.














# 5.7 Other polymers (i)

There are no such residues in this entry.

# 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



# 6 Fit of model and data (i)

### 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	$\# \mathrm{RSRZ}{>}2$	$OWAB(Å^2)$	Q<0.9
1	A	237/237 (100%)	-0.46	0 100 100	11, 17, 30, 43	0
1	В	237/237 (100%)	-0.30	3 (1%) 77 79	11, 18, 39, 54	0
1	С	224/237 (94%)	-0.37	7 (3%) 49 51	10, 17, 40, 62	0
1	D	237/237 (100%)	-0.30	7 (2%) 50 53	11, 19, 40, 81	0
1	E	237/237 (100%)	-0.14	5 (2%) 63 66	11, 20, 40, 71	0
1	F	237/237 (100%)	-0.52	0 100 100	9, 16, 27, 42	0
All	All	1409/1422 (99%)	-0.35	22 (1%) 72 74	9, 18, 37, 81	0

All (22) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	Е	214	ALA	9.2
1	D	209	HIS	5.9
1	Е	212	THR	5.6
1	D	214	ALA	5.4
1	Е	215	ALA	4.2
1	D	207	ARG	3.8
1	С	100	LEU	3.8
1	Е	213	THR	3.7
1	С	97	HIS	3.5
1	D	208	THR	3.4
1	С	98	VAL	2.8
1	Е	211	GLN	2.7
1	С	204	ALA	2.5
1	В	217	ALA	2.3
1	D	212	THR	2.3
1	С	147	ASP	2.3
1	В	212	THR	2.3
1	С	205	HIS	2.3
1	С	220	THR	2.2



Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	213	THR	2.1
1	D	211	GLN	2.1
1	В	97	HIS	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates (i)

There are no monosaccharides in this entry.


## 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	${f B\text{-}factors}({f \AA}^2)$	Q < 0.9
2	FMC	F	301	19/19	0.82	0.20	15,15,15,15	19
2	FMC	С	301	19/19	0.83	0.20	15,15,15,15	19
2	FMC	D	302	19/19	0.85	0.19	15,15,15,15	19
2	FMC	A	300	19/19	0.85	0.24	15,15,15,15	19
4	SO4	С	304	5/5	0.88	0.13	38,38,52,53	5
4	SO4	D	303	5/5	0.92	0.20	36,39,42,44	5
4	SO4	С	303	5/5	0.94	0.15	36,37,43,47	5
4	SO4	Ε	302	5/5	0.96	0.11	29,32,34,43	5
3	PO4	В	301	5/5	0.97	0.07	30,31,36,38	0
4	SO4	F	303	5/5	0.97	0.09	16,20,25,29	5
3	PO4	E	301	5/5	0.98	0.07	24,27,33,33	0
3	PO4	D	301	5/5	0.98	0.08	24,24,28,29	0
3	PO4	F	302	5/5	0.99	0.08	10,10,14,15	0
3	PO4	A	301	5/5	0.99	0.11	13,13,16,16	0
3	PO4	С	302	5/5	0.99	0.05	23,23,27,31	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.














# Electron density around FMC D 302:

 $2 {\rm mF}_o\text{-}{\rm DF}_c$  (at 0.7 rmsd) in gray  ${\rm mF}_o\text{-}{\rm DF}_c$  (at 3 rmsd) in purple (negative) and green (positive)










#### Electron density around FMC A 300:

 $2 \mathrm{mF}_o\text{-DF}_c$  (at 0.7 rmsd) in gray  $\mathrm{mF}_o\text{-DF}_c$  (at 3 rmsd) in purple (negative) and green (positive)









# 6.5 Other polymers (i)

There are no such residues in this entry.

