



# Full wwPDB X-ray Structure Validation Report ⓘ

May 29, 2019 – 04:51 PM EDT

PDB ID : 6NJI  
Title : Crystal Structure of the PDE4D Catalytic Domain and UCR2 Regulatory Helix with T-49  
Authors : Fox III, D.; Fairman, J.W.; Gurney, M.E.  
Deposited on : 2019-01-03  
Resolution : 2.45 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.0 (224370), CSD as540be (2019)  
Xtriage (Phenix) : 1.13  
EDS : rb-20031633  
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)  
Refmac : 5.8.0158  
CCP4 : 7.0 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20031633


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.45 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	111664	1259 (2.48-2.44)
Clashscore	122126	1323 (2.48-2.44)
Ramachandran outliers	120053	1314 (2.48-2.44)
Sidechain outliers	120020	1314 (2.48-2.44)
RSRZ outliers	108989	1238 (2.48-2.44)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	370	 84% 7% • 8%
1	B	370	 85% 5% • 9%

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 5333 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called cAMP-specific 3',5'-cyclic phosphodiesterase 4D.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	339	2639	1674	448	503	14	0	0	0
1	B	335	2571	1629	441	488	13	0	0	0

There are 38 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	243	MET	-	expression tag	UNP Q08499
A	579	ALA	SER	engineered mutation	UNP Q08499
A	581	ALA	SER	engineered mutation	UNP Q08499
A	583	PRO	-	linker	UNP Q08499
A	584	LEU	-	linker	UNP Q08499
A	585	ASP	-	linker	UNP Q08499
A	586	GLU	-	linker	UNP Q08499
A	587	GLN	-	linker	UNP Q08499
A	588	ASN	-	linker	UNP Q08499
A	589	ARG	-	linker	UNP Q08499
A	590	ASP	-	linker	UNP Q08499
A	591	SER	-	linker	UNP Q08499
A	592	GLN	-	linker	UNP Q08499
A	607	GLU	-	expression tag	UNP Q08499
A	608	ASN	-	expression tag	UNP Q08499
A	609	LEU	-	expression tag	UNP Q08499
A	610	TYR	-	expression tag	UNP Q08499
A	611	PHE	-	expression tag	UNP Q08499
A	612	GLN	-	expression tag	UNP Q08499
B	243	MET	-	expression tag	UNP Q08499
B	579	ALA	SER	engineered mutation	UNP Q08499
B	581	ALA	SER	engineered mutation	UNP Q08499
B	583	PRO	-	linker	UNP Q08499
B	584	LEU	-	linker	UNP Q08499
B	585	ASP	-	linker	UNP Q08499

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	586	GLU	-	linker	UNP Q08499
B	587	GLN	-	linker	UNP Q08499
B	588	ASN	-	linker	UNP Q08499
B	589	ARG	-	linker	UNP Q08499
B	590	ASP	-	linker	UNP Q08499
B	591	SER	-	linker	UNP Q08499
B	592	GLN	-	linker	UNP Q08499
B	607	GLU	-	expression tag	UNP Q08499
B	608	ASN	-	expression tag	UNP Q08499
B	609	LEU	-	expression tag	UNP Q08499
B	610	TYR	-	expression tag	UNP Q08499
B	611	PHE	-	expression tag	UNP Q08499
B	612	GLN	-	expression tag	UNP Q08499


- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	1	Total Zn 1 1	0	0
2	A	1	Total Zn 1 1	0	0

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	1	Total Mg 1 1	0	0
3	A	1	Total Mg 1 1	0	0

- Molecule 4 is 2-(4-{[4-(3-chlorophenyl)-6-ethyl-1,3,5-triazin-2-yl]amino}phenyl)ethan-1-ol (three-letter code: KR4) (formula: C<sub>19</sub>H<sub>19</sub>ClN<sub>4</sub>O).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
			Total	C	Cl	N			O
4	A	1	25	19	1	4	1	0	0
4	B	1	25	19	1	4	1	0	0

- Molecule 5 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
5	A	46	46	46	0	0
5	B	23	23	23	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: cAMP-specific 3',5'-cyclic phosphodiesterase 4D

Chain A: 84% 7% 8%



- Molecule 1: cAMP-specific 3',5'-cyclic phosphodiesterase 4D

Chain B: 85% 5% 9%



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	124.04Å 85.66Å 83.45Å 90.00° 110.41° 90.00°	Depositor
Resolution (Å)	42.87 – 2.45 42.83 – 2.45	Depositor EDS
% Data completeness (in resolution range)	99.0 (42.87-2.45) 98.1 (42.83-2.45)	Depositor EDS
$R_{merge}$	0.05	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.52 (at 2.45Å)	Xtrriage
Refinement program	REFMAC 5.7.0029	Depositor
R, $R_{free}$	0.184 , 0.230 0.188 , 0.223	Depositor DCC
$R_{free}$ test set	1504 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	47.2	Xtrriage
Anisotropy	0.084	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 39.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	5333	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	53.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.71% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, MG, KR4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.76	0/2694	0.78	0/3679
1	B	0.66	0/2625	0.73	0/3589
All	All	0.71	0/5319	0.76	0/7268

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2639	0	2497	14	0
1	B	2571	0	2377	11	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	25	0	0	1	0
4	B	25	0	0	1	0
5	A	46	0	0	1	0
5	B	23	0	0	0	0
All	All	5333	0	4874	27	0



The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (27) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:453:VAL:O	1:A:456:LYS:HG2	1.96	0.66
1:A:489:THR:HB	1:A:561:ILE:HG23	1.88	0.56
1:A:298:LEU:HD23	1:A:303:ILE:O	2.08	0.53
1:B:298:LEU:HD23	1:B:303:ILE:O	2.11	0.51
1:B:282:ARG:N	1:B:283:PRO:CD	2.77	0.48
1:A:409:GLU:HB2	1:A:412:CYS:SG	2.54	0.48
1:A:487:ASN:HB2	1:A:488:PRO:HD3	1.96	0.48
1:A:523:MET:CE	1:A:596:VAL:HA	2.44	0.47
1:A:555:HIS:HA	1:A:556:PRO:HA	1.62	0.47
1:B:523:MET:CE	1:B:596:VAL:HA	2.45	0.47
1:A:253:GLU:O	1:A:257:VAL:HG23	2.15	0.47
1:B:487:ASN:HB2	1:B:488:PRO:HD3	1.97	0.47
4:B:702:KR4:C10	4:B:702:KR4:N1	2.77	0.45
4:A:702:KR4:N1	4:A:702:KR4:C10	2.80	0.45
1:B:409:GLU:HB2	1:B:412:CYS:SG	2.57	0.45
1:B:555:HIS:HA	1:B:556:PRO:HA	1.66	0.45
1:A:295:ARG:HE	1:A:339:HIS:CE1	2.36	0.44
1:A:282:ARG:N	1:A:283:PRO:CD	2.80	0.44
1:B:557:ASP:N	1:B:557:ASP:OD1	2.51	0.44
1:A:303:ILE:O	1:A:304:PRO:C	2.54	0.43
1:B:389:TYR:CE1	1:B:397:ASN:HB3	2.54	0.42
1:A:441:LYS:HE3	5:A:825:HOH:O	2.20	0.41
1:A:429:MET:O	1:A:433:ILE:HG13	2.20	0.41
1:A:350:VAL:HG11	1:A:466:LEU:HD12	2.03	0.41
1:B:328:ASN:OD1	1:B:328:ASN:N	2.54	0.41
1:B:429:MET:O	1:B:433:ILE:HG13	2.21	0.41
1:B:367:ASP:O	1:B:370:HIS:HB2	2.20	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	335/370 (90%)	324 (97%)	10 (3%)	1 (0%)	43	52
1	B	329/370 (89%)	316 (96%)	12 (4%)	1 (0%)	43	52
All	All	664/740 (90%)	640 (96%)	22 (3%)	2 (0%)	43	52

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	515	GLU
1	B	515	GLU

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	284/338 (84%)	280 (99%)	4 (1%)	69	79
1	B	267/338 (79%)	264 (99%)	3 (1%)	76	83
All	All	551/676 (82%)	544 (99%)	7 (1%)	71	81

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	423	ARG
1	A	426	LEU
1	A	456	LYS
1	A	556	PRO
1	B	423	ARG
1	B	556	PRO
1	B	557	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	535	GLN
1	B	535	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 4 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	KR4	A	702	-	26,27,27	1.20	5 (19%)	36,36,36	2.29	7 (19%)
4	KR4	B	702	-	26,27,27	1.13	2 (7%)	36,36,36	2.64	11 (30%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	KR4	A	702	-	-	0/13/13/13	0/3/3/3
4	KR4	B	702	-	-	0/13/13/13	0/3/3/3

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	702	KR4	C5-N3	-2.76	1.34	1.40
4	A	702	KR4	C3-N1	2.06	1.38	1.34
4	A	702	KR4	C2-N	2.11	1.36	1.33
4	A	702	KR4	C2-N2	2.16	1.36	1.33
4	A	702	KR4	C17-CL	2.50	1.80	1.74
4	B	702	KR4	C2-N2	2.55	1.37	1.33
4	B	702	KR4	C17-CL	2.96	1.81	1.74

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	702	KR4	N1-C4-N2	-5.47	117.71	126.22
4	B	702	KR4	N1-C4-N2	-5.40	117.83	126.22
4	B	702	KR4	N-C2-N2	-4.28	120.41	126.06
4	A	702	KR4	N-C2-N2	-4.21	120.50	126.06
4	B	702	KR4	C5-N3-C4	-3.28	120.97	129.31
4	A	702	KR4	C5-N3-C4	-3.26	121.02	129.31
4	B	702	KR4	N-C3-N1	-2.94	120.68	125.28
4	B	702	KR4	C13-C18-C17	2.03	121.67	119.14
4	B	702	KR4	N3-C4-N1	2.40	124.95	116.90
4	A	702	KR4	N3-C4-N1	2.71	125.98	116.90
4	B	702	KR4	C1-C2-N	2.80	121.27	117.29
4	B	702	KR4	C13-C3-N1	3.34	122.89	117.31
4	A	702	KR4	C1-C2-N	3.64	122.47	117.29
4	B	702	KR4	C2-N-C3	4.71	118.64	114.80
4	B	702	KR4	C4-N2-C2	5.04	121.73	114.54
4	A	702	KR4	C4-N2-C2	5.80	122.82	114.54
4	A	702	KR4	C4-N1-C3	7.14	119.15	114.60
4	B	702	KR4	C4-N1-C3	9.41	120.60	114.60

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	702	KR4	1	0
4	B	702	KR4	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	339/370 (91%)	-0.44	1 (0%) 93 95	29, 47, 68, 96	0
1	B	335/370 (90%)	-0.26	2 (0%) 89 90	32, 57, 86, 102	0
All	All	674/740 (91%)	-0.35	3 (0%) 92 93	29, 51, 81, 102	0

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	258	LEU	2.8
1	B	463	VAL	2.6
1	B	347	LEU	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	KR4	B	702	25/25	0.95	0.24	50,55,62,73	0
4	KR4	A	702	25/25	0.96	0.20	40,45,53,56	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	MG	A	701	1/1	0.97	0.08	36,36,36,36	0
3	MG	B	701	1/1	0.97	0.12	42,42,42,42	0
2	ZN	B	700	1/1	1.00	0.14	44,44,44,44	0
2	ZN	A	700	1/1	1.00	0.15	38,38,38,38	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.