



# Full wwPDB X-ray Structure Validation Report ⓘ

Jan 14, 2024 – 01:03 am GMT

PDB ID : 6QD5  
Title : X-ray Structure of the Human Urea Channel SLC14A1/UT1  
Authors : Dietz, L.; Chi, G.; Pike, A.C.W.; Moreau, C.; Man, H.; Snee, M.; Scacioc, A.; Shrestha, L.; Mukhopadhyay, S.M.M.; Mckinley, G.; Ellis, K.; Kliszczak, M.; Chalk, R.; Borkowska, O.; Burgess-Brown, N.A.; von Delft, F.; Arrow-smith, C.H.; Edwards, A.M.; Bountra, C.; Durr, K.L.; Structural Genomics Consortium (SGC)  
Deposited on : 2018-12-31  
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3907 (2.40-2.40)
Clashscore	141614	4398 (2.40-2.40)
Ramachandran outliers	138981	4318 (2.40-2.40)
Sidechain outliers	138945	4319 (2.40-2.40)
RSRZ outliers	127900	3811 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	367	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
 Validation Pipeline (wwPDB-VP) : 2.36

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	BOG	A	405	-	-	-	X
2	BOG	A	407	-	-	-	X

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 2974 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Urea transporter 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	356	2702	1795	417	466	24	0	0	0

There are 11 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	30	MET	-	initiating methionine	UNP Q13336
A	167	VAL	MET	variant	UNP Q13336
A	211	GLN	ASN	engineered mutation	UNP Q13336
A	280	ASN	ASP	variant	UNP Q13336
A	390	ALA	-	expression tag	UNP Q13336
A	391	GLU	-	expression tag	UNP Q13336
A	392	ASN	-	expression tag	UNP Q13336
A	393	LEU	-	expression tag	UNP Q13336
A	394	TYR	-	expression tag	UNP Q13336
A	395	PHE	-	expression tag	UNP Q13336
A	396	GLN	-	expression tag	UNP Q13336

- Molecule 2 is octyl beta-D-glucopyranoside (three-letter code: BOG) (formula: C<sub>14</sub>H<sub>28</sub>O<sub>6</sub>).




Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	C O	6	0
			20	14 6		
2	A	1	Total	C O	0	0
			20	14 6		
2	A	1	Total	C O	0	0
			20	14 6		
2	A	1	Total	C O	7	0
			20	14 6		
2	A	1	Total	C O	0	0
			20	14 6		
2	A	1	Total	C O	5	0
			20	14 6		
2	A	1	Total	C O	2	0
			20	14 6		

- Molecule 3 is CHOLESTEROL HEMISUCCINATE (three-letter code: Y01) (formula:  $C_{31}H_{50}O_4$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			35	31	4		

- Molecule 4 is TETRAETHYLENE GLYCOL (three-letter code: PG4) (formula:  $C_8H_{18}O_5$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			13	8	5		
4	A	1	Total	C	O	0	0
			13	8	5		


- Molecule 5 is water.

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
5	A	71	Total	O	0	0
			71	71		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Urea transporter 1





## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 63	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	112.33Å 112.33Å 87.94Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	47.34 – 2.40 47.33 – 2.40	Depositor EDS
% Data completeness (in resolution range)	99.7 (47.34-2.40) 99.7 (47.33-2.40)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.49 (at 2.39Å)	Xtrriage
Refinement program	PHENIX (1.12_2829: ???)	Depositor
R, $R_{free}$	0.175 , 0.225 0.175 , 0.225	Depositor DCC
$R_{free}$ test set	1165 reflections (4.70%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	50.8	Xtrriage
Anisotropy	0.113	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 60.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.065 for h,-h-k,-l	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	2974	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	61.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.39% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BOG, Y01, PG4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.45	0/2779	0.56	0/3801

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2702	0	2737	18	0
2	A	140	0	196	1	0
3	A	35	0	49	0	0
4	A	26	0	36	0	0
5	A	71	0	0	1	0
All	All	2974	0	3018	18	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (18) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:212:ILE:HD12	1:A:214:TRP:HE1	1.55	0.71
1:A:41:ASP:OD1	1:A:43:LYS:HE2	1.98	0.63
1:A:148:LEU:HD23	2:A:403:BOG:H7'2	1.82	0.60
1:A:31:PHE:O	1:A:35:LEU:HB2	2.04	0.56
1:A:193:HIS:HD2	1:A:200:ALA:O	1.89	0.55
1:A:106:MET:HB3	1:A:158:PRO:HG2	1.91	0.52
1:A:106:MET:HE3	1:A:110:LEU:HG	1.95	0.49
1:A:71:PHE:HZ	1:A:342:PHE:CZ	2.31	0.48
1:A:61:TRP:CD1	1:A:118:ALA:HA	2.49	0.48
1:A:279:GLU:H	1:A:279:GLU:CD	2.17	0.47
1:A:361:LYS:NZ	5:A:501:HOH:O	2.48	0.46
1:A:131:VAL:O	1:A:135:VAL:HG23	2.16	0.45
1:A:366:LYS:HA	1:A:366:LYS:HD3	1.86	0.43
1:A:217:LEU:HD11	1:A:222:LEU:HD11	2.00	0.43
1:A:106:MET:CE	1:A:110:LEU:HG	2.48	0.42
1:A:224:LYS:O	1:A:228:VAL:HG23	2.20	0.41
1:A:323:GLY:HA2	1:A:341:PRO:CG	2.51	0.41
1:A:151:CYS:O	1:A:155:MET:HG3	2.21	0.40

There are no symmetry-related clashes.

### 5.3 Torsion angles [i](#)

#### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	354/367 (96%)	345 (98%)	9 (2%)	0	<b>100</b> <b>100</b>

There are no Ramachandran outliers to report.

#### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	288/306 (94%)	286 (99%)	2 (1%)	84 92

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	43	LYS
1	A	51	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	193	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

10 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	BOG	A	404	-	20,20,20	0.13	0	25,25,25	0.19	0
4	PG4	A	409	-	12,12,12	0.50	0	11,11,11	0.54	0
2	BOG	A	403	-	20,20,20	0.21	0	25,25,25	0.33	0
2	BOG	A	405	-	20,20,20	0.32	0	25,25,25	0.51	0
4	PG4	A	410	-	12,12,12	0.50	0	11,11,11	0.50	0
2	BOG	A	407	-	20,20,20	0.12	0	25,25,25	0.33	0
3	Y01	A	408	-	38,38,38	1.42	6 (15%)	57,57,57	1.71	13 (22%)
2	BOG	A	402	-	20,20,20	0.23	0	25,25,25	0.40	0
2	BOG	A	406	-	20,20,20	0.15	0	25,25,25	0.33	0
2	BOG	A	401	-	20,20,20	0.23	0	25,25,25	0.50	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BOG	A	404	-	-	8/11/31/31	0/1/1/1
4	PG4	A	409	-	-	6/10/10/10	-
2	BOG	A	403	-	-	0/11/31/31	0/1/1/1
2	BOG	A	405	-	-	5/11/31/31	0/1/1/1
4	PG4	A	410	-	-	7/10/10/10	-
2	BOG	A	407	-	-	4/11/31/31	0/1/1/1
3	Y01	A	408	-	-	3/19/77/77	0/4/4/4
2	BOG	A	402	-	-	4/11/31/31	0/1/1/1
2	BOG	A	406	-	-	3/11/31/31	0/1/1/1
2	BOG	A	401	-	-	7/11/31/31	0/1/1/1

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	408	Y01	CAK-CAI	3.11	1.56	1.50
3	A	408	Y01	CAQ-CAP	2.83	1.61	1.54
3	A	408	Y01	CBI-CBE	-2.79	1.49	1.55
3	A	408	Y01	OAW-CBC	-2.70	1.39	1.46
3	A	408	Y01	OAW-CAY	2.57	1.41	1.34
3	A	408	Y01	CAS-CBF	-2.46	1.49	1.53

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	408	Y01	OAW-CAY-CAM	4.54	121.28	111.50
3	A	408	Y01	CAK-CAI-CAZ	-4.37	117.00	125.06
3	A	408	Y01	CAP-CAQ-CBG	-3.87	97.45	105.13
3	A	408	Y01	CBF-CBD-CBG	-3.15	104.87	109.09
3	A	408	Y01	CAK-CBD-CBG	-3.12	106.38	110.91
3	A	408	Y01	CAS-CBF-CBH	-3.00	109.13	113.08
3	A	408	Y01	CBI-CBG-CBD	-2.74	110.32	114.38
3	A	408	Y01	CAP-CBE-CBB	-2.39	108.45	112.15
3	A	408	Y01	CBC-OAW-CAY	-2.38	111.94	117.79
3	A	408	Y01	CAO-CBB-CBE	-2.35	105.43	110.28
3	A	408	Y01	CAD-CBH-CBF	-2.29	108.95	111.68
3	A	408	Y01	OAW-CAY-OAG	-2.28	118.20	123.70
3	A	408	Y01	CAQ-CBG-CBI	2.02	106.28	103.84

There are no chirality outliers.

All (47) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	401	BOG	O5-C1-O1-C1'
2	A	402	BOG	C2'-C1'-O1-C1
2	A	405	BOG	C2'-C1'-O1-C1
4	A	409	PG4	C3-C4-O3-C5
2	A	404	BOG	O5-C5-C6-O6
4	A	409	PG4	O3-C5-C6-O4
4	A	409	PG4	C5-C6-O4-C7
2	A	401	BOG	C2-C1-O1-C1'
2	A	405	BOG	C2-C1-O1-C1'
2	A	404	BOG	C4-C5-C6-O6
2	A	407	BOG	O5-C5-C6-O6
2	A	404	BOG	O1-C1'-C2'-C3'
2	A	404	BOG	C2-C1-O1-C1'
4	A	410	PG4	O1-C1-C2-O2
2	A	401	BOG	C2'-C1'-O1-C1
2	A	405	BOG	C3'-C4'-C5'-C6'
2	A	401	BOG	C2'-C3'-C4'-C5'
2	A	404	BOG	O5-C1-O1-C1'
2	A	401	BOG	O5-C5-C6-O6
4	A	409	PG4	O1-C1-C2-O2
2	A	405	BOG	O5-C5-C6-O6
2	A	404	BOG	C3'-C4'-C5'-C6'
2	A	404	BOG	C2'-C1'-O1-C1
2	A	407	BOG	C2'-C1'-O1-C1
2	A	404	BOG	C1'-C2'-C3'-C4'

*Continued on next page...*

*Continued from previous page...*




Mol	Chain	Res	Type	Atoms
4	A	410	PG4	C3-C4-O3-C5
2	A	402	BOG	C4-C5-C6-O6
4	A	410	PG4	C4-C3-O2-C2
2	A	406	BOG	C5'-C6'-C7'-C8'
2	A	406	BOG	C2'-C1'-O1-C1
4	A	410	PG4	O4-C7-C8-O5
2	A	402	BOG	C1'-C2'-C3'-C4'
4	A	409	PG4	C4-C3-O2-C2
2	A	401	BOG	C1'-C2'-C3'-C4'
2	A	407	BOG	C2'-C3'-C4'-C5'
2	A	405	BOG	C2'-C3'-C4'-C5'
2	A	401	BOG	C3'-C4'-C5'-C6'
4	A	410	PG4	C5-C6-O4-C7
3	A	408	Y01	CAM-CAL-CAX-OAF
3	A	408	Y01	CAM-CAL-CAX-OAH
2	A	402	BOG	C3'-C4'-C5'-C6'
4	A	410	PG4	O2-C3-C4-O3
4	A	409	PG4	O2-C3-C4-O3
2	A	406	BOG	C4'-C5'-C6'-C7'
2	A	407	BOG	C4-C5-C6-O6
3	A	408	Y01	CAL-CAM-CAY-OAW
4	A	410	PG4	O3-C5-C6-O4

There are no ring outliers.


1 monomer is involved in 1 short contact:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	403	BOG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	356/367 (97%)	0.15	27 (7%) <b>13</b> <b>12</b>	38, 54, 97, 150	0

All (27) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	32	PRO	9.2
1	A	35	LEU	8.3
1	A	33	LYS	7.8
1	A	384	MET	6.2
1	A	34	ALA	6.1
1	A	31	PHE	5.8
1	A	217	LEU	5.5
1	A	37	TYR	5.4
1	A	36	GLY	5.2
1	A	385	VAL	4.9
1	A	383	ARG	4.8
1	A	171	TRP	4.7
1	A	38	VAL	4.5
1	A	215	SER	4.4
1	A	216	ASP	4.0
1	A	220	LEU	3.8
1	A	214	TRP	3.3
1	A	213	SER	3.3
1	A	223	LEU	3.1
1	A	218	SER	2.9
1	A	219	ALA	2.9
1	A	212	ILE	2.7
1	A	386	GLU	2.7
1	A	207	THR	2.5
1	A	39	THR	2.5
1	A	211	GLN	2.3
1	A	306	TRP	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	BOG	A	405	20/20	0.65	0.46	88,119,139,140	0
2	BOG	A	407	20/20	0.73	0.42	71,113,132,140	2
2	BOG	A	401	20/20	0.80	0.26	69,100,112,118	6
4	PG4	A	409	13/13	0.84	0.19	59,74,88,93	0
2	BOG	A	404	20/20	0.87	0.35	126,131,142,143	7
2	BOG	A	403	20/20	0.87	0.24	71,89,100,109	0
4	PG4	A	410	13/13	0.87	0.18	62,70,102,103	0
3	Y01	A	408	35/35	0.88	0.15	50,67,90,100	0
2	BOG	A	406	20/20	0.90	0.17	82,98,122,125	5
2	BOG	A	402	20/20	0.94	0.12	55,59,65,70	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around BOG A 405:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around BOG A 407:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around BOG A 401:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around BOG A 404:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)







**Electron density around Y01 A 408:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around BOG A 406:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [i](#)

There are no such residues in this entry.