



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 23, 2024 – 10:28 AM EDT

PDB ID : 6SQV  
Title : Structure of the U1A variant A1-98 Y31H/Q36R/R70W  
Authors : Rosenbach, H.; Span, I.  
Deposited on : 2019-09-04  
Resolution : 2.45 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 1.20.1  
EDS : 2.37.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.37.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.45 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1544 (2.48-2.44)
Clashscore	141614	1613 (2.48-2.44)
Ramachandran outliers	138981	1598 (2.48-2.44)
Sidechain outliers	138945	1598 (2.48-2.44)
RSRZ outliers	127900	1523 (2.48-2.44)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	AAA	98	 7% 73% 15% • 9%
1	BBB	98	 3% 72% 14% 13%
1	CCC	98	 7% 72% 10% 5% 12%
1	DDD	98	 4% 72% 10% • 16%

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 5718 atoms, of which 2892 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called U1 small nuclear ribonucleoprotein A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
1	AAA	89	1478	470	749	127	129	3	37	0	0
1	BBB	85	1419	451	720	123	122	3	36	0	0
1	CCC	86	1429	456	725	123	122	3	35	0	0
1	DDD	82	1372	438	698	117	116	3	35	0	0

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
AAA	31	HIS	TYR	engineered mutation	UNP P09012
AAA	36	ARG	GLN	engineered mutation	UNP P09012
AAA	70	TRP	ARG	engineered mutation	UNP P09012
AAA	98	TRP	-	expression tag	UNP P09012
BBB	31	HIS	TYR	engineered mutation	UNP P09012
BBB	36	ARG	GLN	engineered mutation	UNP P09012
BBB	70	TRP	ARG	engineered mutation	UNP P09012
BBB	98	TRP	-	expression tag	UNP P09012
CCC	31	HIS	TYR	engineered mutation	UNP P09012
CCC	36	ARG	GLN	engineered mutation	UNP P09012
CCC	70	TRP	ARG	engineered mutation	UNP P09012
CCC	98	TRP	-	expression tag	UNP P09012
DDD	31	HIS	TYR	engineered mutation	UNP P09012
DDD	36	ARG	GLN	engineered mutation	UNP P09012
DDD	70	TRP	ARG	engineered mutation	UNP P09012
DDD	98	TRP	-	expression tag	UNP P09012

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	AAA	1	Total O S 5 4 1	0	0
2	AAA	1	Total O S 5 4 1	0	0
2	CCC	1	Total O S 5 4 1	0	0



- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	BBB	1	Total O 1 1	0	0
3	CCC	1	Total O 1 1	0	0
3	DDD	3	Total O 3 3	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: U1 small nuclear ribonucleoprotein A




- Molecule 1: U1 small nuclear ribonucleoprotein A



- Molecule 1: U1 small nuclear ribonucleoprotein A



- Molecule 1: U1 small nuclear ribonucleoprotein A



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	76.23Å 76.43Å 137.35Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	51.08 – 2.45 51.02 – 2.45	Depositor EDS
% Data completeness (in resolution range)	99.9 (51.08-2.45) 99.9 (51.02-2.45)	Depositor EDS
$R_{merge}$	0.05	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.94 (at 2.45Å)	Xtrriage
Refinement program	REFMAC 5.8.0253	Depositor
R, $R_{free}$	0.240 , 0.273 0.243 , 0.274	Depositor DCC
$R_{free}$ test set	1497 reflections (4.96%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	79.2	Xtrriage
Anisotropy	0.016	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 54.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.016 for k,h,-l	Xtrriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	5718	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	83.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.06% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	AAA	0.72	0/745	0.90	0/1001
1	BBB	0.70	0/714	0.87	0/957
1	CCC	0.74	0/720	0.95	0/968
1	DDD	0.65	0/689	0.86	0/923
All	All	0.70	0/2868	0.90	0/3849

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	AAA	729	749	746	12	0
1	BBB	699	720	717	12	0
1	CCC	704	725	722	10	0
1	DDD	674	698	696	4	0
2	AAA	10	0	0	0	0
2	CCC	5	0	0	0	0
3	BBB	1	0	0	0	0
3	CCC	1	0	0	0	0
3	DDD	3	0	0	0	0
All	All	2826	2892	2881	35	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (35) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:AAA:14:ILE:HG21	1:AAA:82:MET:HE3	1.61	0.83
1:AAA:14:ILE:HG21	1:AAA:82:MET:CE	2.12	0.80
1:BBB:7:ARG:NH1	1:BBB:7:ARG:HB2	2.06	0.70
1:AAA:14:ILE:HD13	1:AAA:82:MET:HE1	1.81	0.61
1:CCC:29:SER:O	1:CCC:32:ALA:HB3	2.01	0.61
1:BBB:7:ARG:N	1:BBB:8:PRO:CD	2.66	0.58
1:CCC:49:LEU:C	1:CCC:51:MET:H	2.05	0.56
1:CCC:13:TYR:OH	1:CCC:54:GLN:NE2	2.40	0.55
1:BBB:26:LEU:HD23	1:BBB:45:VAL:HG11	1.91	0.53
1:BBB:69:LEU:HD23	1:BBB:69:LEU:O	2.09	0.53
1:BBB:10:HIS:HD2	1:BBB:59:PHE:O	1.93	0.52
1:BBB:69:LEU:HD23	1:BBB:69:LEU:C	2.31	0.51
1:CCC:50:LYS:O	1:CCC:50:LYS:HD2	2.10	0.51
1:DDD:31:HIS:O	1:DDD:35:SER:HB3	2.11	0.51
1:CCC:6:THR:O	1:CCC:8:PRO:HD3	2.11	0.50
1:AAA:14:ILE:CG2	1:AAA:82:MET:CE	2.87	0.49
1:BBB:7:ARG:HB2	1:BBB:7:ARG:CZ	2.42	0.49
1:AAA:14:ILE:CG2	1:AAA:82:MET:HE3	2.38	0.48
1:BBB:10:HIS:CD2	1:BBB:59:PHE:O	2.68	0.47
1:AAA:17:LEU:HD21	1:AAA:82:MET:HE3	1.99	0.45
1:AAA:4:PRO:HG3	1:BBB:69:LEU:HD22	2.00	0.44
1:AAA:14:ILE:HD13	1:AAA:82:MET:CE	2.46	0.44
1:AAA:13:TYR:OH	1:AAA:54:GLN:HG2	2.17	0.44
1:AAA:23:LYS:HG2	1:AAA:45:VAL:HG23	1.99	0.43
1:BBB:25:GLU:OE1	1:CCC:28:LYS:HD3	2.19	0.43
1:AAA:6:THR:O	1:AAA:8:PRO:HD3	2.19	0.43
1:CCC:52:ARG:CG	1:CCC:52:ARG:HH21	2.32	0.42
1:AAA:78:TYR:CZ	1:DDD:32:ALA:HB2	2.55	0.42
1:CCC:48:SER:OG	1:CCC:49:LEU:N	2.52	0.42
1:CCC:49:LEU:C	1:CCC:51:MET:N	2.72	0.41
1:DDD:51:MET:HA	1:DDD:54:GLN:NE2	2.35	0.41
1:BBB:7:ARG:N	1:BBB:8:PRO:HD2	2.36	0.41
1:CCC:49:LEU:O	1:CCC:51:MET:N	2.53	0.41
1:DDD:9:ASN:OD1	1:DDD:10:HIS:N	2.54	0.41
1:BBB:11:THR:HG22	1:BBB:87:ALA:HB2	2.03	0.40

There are no symmetry-related clashes.



## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	AAA	87/98 (89%)	82 (94%)	5 (6%)	0	100	100
1	BBB	83/98 (85%)	81 (98%)	2 (2%)	0	100	100
1	CCC	84/98 (86%)	79 (94%)	3 (4%)	2 (2%)	6	3
1	DDD	80/98 (82%)	78 (98%)	2 (2%)	0	100	100
All	All	334/392 (85%)	320 (96%)	12 (4%)	2 (1%)	25	29

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	CCC	48	SER
1	CCC	50	LYS

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	AAA	81/88 (92%)	73 (90%)	8 (10%)	8	7
1	BBB	77/88 (88%)	73 (95%)	4 (5%)	23	30
1	CCC	77/88 (88%)	71 (92%)	6 (8%)	12	15
1	DDD	74/88 (84%)	69 (93%)	5 (7%)	16	19
All	All	309/352 (88%)	286 (93%)	23 (7%)	13	16

All (23) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	AAA	3	VAL
1	AAA	23	LYS
1	AAA	47	ARG
1	AAA	52	ARG
1	AAA	62	VAL
1	AAA	82	MET
1	AAA	89	THR
1	AAA	91	SER
1	BBB	9	ASN
1	BBB	52	ARG
1	BBB	88	LYS
1	BBB	90	ASP
1	CCC	24	ASP
1	CCC	46	SER
1	CCC	47	ARG
1	CCC	49	LEU
1	CCC	52	ARG
1	CCC	54	GLN
1	DDD	24	ASP
1	DDD	27	LYS
1	DDD	35	SER
1	DDD	36	ARG
1	DDD	46	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	SO4	AAA	101	-	4,4,4	0.24	0	6,6,6	0.12	0
2	SO4	AAA	102	-	4,4,4	0.25	0	6,6,6	0.21	0
2	SO4	CCC	101	-	4,4,4	0.30	0	6,6,6	0.13	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	AAA	89/98 (90%)	0.69	7 (7%) 12 9	54, 72, 124, 163	0
1	BBB	85/98 (86%)	0.53	3 (3%) 44 40	58, 75, 127, 158	0
1	CCC	86/98 (87%)	0.61	7 (8%) 12 9	52, 77, 123, 153	0
1	DDD	82/98 (83%)	0.66	4 (4%) 29 27	58, 75, 127, 159	0
All	All	342/392 (87%)	0.62	21 (6%) 21 18	52, 75, 130, 163	0

All (21) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	BBB	49	LEU	5.3
1	DDD	52	ARG	5.3
1	AAA	49	LEU	5.2
1	AAA	47	ARG	5.1
1	DDD	51	MET	5.1
1	CCC	51	MET	4.8
1	AAA	52	ARG	4.7
1	CCC	49	LEU	4.5
1	AAA	23	LYS	4.3
1	AAA	45	VAL	3.8
1	BBB	51	MET	3.4
1	CCC	48	SER	3.0
1	AAA	50	LYS	2.9
1	CCC	45	VAL	2.7
1	CCC	52	ARG	2.7
1	DDD	50	LYS	2.6
1	CCC	47	ARG	2.4
1	DDD	49	LEU	2.4
1	AAA	51	MET	2.3
1	BBB	50	LYS	2.3
1	CCC	50	LYS	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	SO4	AAA	101	5/5	0.93	0.10	99,103,110,134	0
2	SO4	AAA	102	5/5	0.93	0.12	108,114,121,135	0
2	SO4	CCC	101	5/5	0.94	0.17	102,114,127,133	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.